

The St Petersburg Paradox (and inductive logic)

Selmer Bringsjord

Rensselaer AI & Reasoning (RAIR) Lab
Department of Cognitive Science
Department of Computer Science
Lally School of Management & Technology
Rensselaer Polytechnic Institute (RPI)
Troy, New York 12180 USA

IFLAI2 2022
10/17/2022




R&D enabled by:



The St Petersburg Paradox ...

Ignore Those Who Say WWWTBAM is an Instance



Ignore Those Who Say WWWTBAM is an Instance

Presently rational within this game!



Here's the game; \$30 to play.



Here's the game; \$30 to play.



Here's the game; \$30 to play.



Here's the game; \$30 to play.



Here's the game; \$30 to play.



\$2

Here's the game; \$30 to play.



\$2



Here's the game; \$30 to play.



\$2



\$4

Here's the game; \$30 to play.



\$2



\$4



Here's the game; \$30 to play.



\$2



\$4



\$8

Here's the game; \$30 to play.



\$2



\$4



\$8



Here's the game; \$30 to play.



\$2



\$4



\$8



\$16

Here's the game; \$30 to play.



\$2



\$4



\$8



\$16

...

For $n = 1, 2, \dots, 10$

n	$prob(n)$	Prize	Expected Payoff
1	1/2	\$2	\$1
2	1/4	\$4	\$1
3	1/8	\$8	\$1
4	1/16	\$16	\$1
5	1/32	\$32	\$1
6	1/64	\$64	\$1
7	1/128	\$128	\$1
8	1/256	\$256	\$1
9	1/512	\$512	\$1
10	1/1024	\$1,024	\$1
...

For $n = 1, 2, \dots, 10$

n	$prob(n)$	Prize	Expected Payoff
1	1/2	\$2	\$1
2	1/4	\$4	\$1
3	1/8	\$8	\$1
4	1/16	\$16	\$1
5	1/32	\$32	\$1
6	1/64	\$64	\$1
7	1/128	\$128	\$1
8	1/256	\$256	\$1
9	1/512	\$512	\$1
10	1/1024	\$1,024	\$1
...	9.5367431640625e-7

For $n = 1, 2, \dots, 10$

n	$\text{prob}(n)$	Prize	Expected Payoff
1	1/2	\$2	\$1
2	1/4	\$4	\$1
3	1/8	\$8	\$1
4	1/16	\$16	\$1
5	1/32	\$32	\$1
6	1/64	\$64	\$1
7	1/128	\$128	\$1
8	1/256	\$256	\$1
9	1/512	\$512	\$1
10	1/1024	\$1,024	\$1
...	9.5367431640625e-7

Theorem: You should give Selmer all your savings ($\$k$) to play the game.

For $n = 1, 2, \dots, 10$

n	$\text{prob}(n)$	Prize	Expected Payoff
1	1/2	\$2	\$1
2	1/4	\$4	\$1
3	1/8	\$8	\$1
4	1/16	\$16	\$1
5	1/32	\$32	\$1
6	1/64	\$64	\$1
7	1/128	\$128	\$1
8	1/256	\$256	\$1
9	1/512	\$512	\$1
10	1/1024	\$1,024	\$1
...	9.5367431640625e-7

Theorem: You should give Selmer all your savings ($\$k$) to play the game.

Proof: The expected value of playing the game is $\$1$. We know that you're rational, so since any finite amount of dollars isn't an infinite amount of dollars, you will pay $\$k$ to us to play. **QED**

The Optimality Principle

When choosing between alternative actions a_1 and a_2 , rationality dictates choosing that action that maximizes expected value, computed by multiplying the value of each outcome that can result from each action by the probability that it will occur, adding the results together, and selecting the action associated with the higher utility.

The Optimality Principle

Bernoulli's Bad Idea

n	$prob(n)$	Prize	Utiles	Expected Utility
1	1/2	\$2	0.301	0.1505
2	1/4	\$4	0.602	0.1505
3	1/8	\$8	0.903	0.1129
4	1/16	\$16	1.204	0.0753
5	1/32	\$32	1.505	0.0470
6	1/64	\$64	1.806	0.0282
7	1/128	\$128	2.107	0.0165
8	1/256	\$256	2.408	0.0094
9	1/512	\$512	2.709	0.0053
10	1/1024	\$1,024	3.010	0.0029
...

Refutation: Selmer & Naveen offer a variant game, based on stacked exponentiation. E.g., let's play a game in which the prize money is 10^{2^n} . Sorry Bernoulli! **QED**

Bernoulli's Bad Idea

n	Probability	Prize Money	Expected Utility	Actual Utility
1	1/1	\$1	0.602	0.602
2	1/2	\$2	0.708	0.505
3	1/4	\$4	0.807	0.354
4	1/8	\$8	0.903	0.253
5	1/16	\$16	1.004	0.176
6	1/32	\$32	1.105	0.112
7	1/64	\$64	1.206	0.075
8	1/128	\$128	1.307	0.050
9	1/256	\$256	1.408	0.033
10	1/512	\$512	1.509	0.022
...

Principle of Decreasing Marginal Utility (DMU): The utility of \$ $k = \log_{10}(k)$.

Refutation: Selmer & Naveen offer a variant game, based on stacked exponentiation. E.g., let's play a game in which the prize money is 10^{2^n} . Sorry Bernoulli! **QED**

Bernoulli's Bad Idea

n	$prob(n)$	Prize	Utiles	Expected Utility
1	1/2	\$2	0.301	0.1505
2	1/4	\$4	0.602	0.1505
3	1/8	\$8	0.903	0.1129
4	1/16	\$16	1.204	0.0753
5	1/32	\$32	1.505	0.0470
6	1/64	\$64	1.806	0.0282
7	1/128	\$128	2.107	0.0165
8	1/256	\$256	2.408	0.0094
9	1/512	\$512	2.709	0.0053
10	1/1024	\$1,024	3.010	0.0029
...

Refutation: Selmer & Naveen offer a variant game, based on stacked exponentiation. E.g., let's play a game in which the prize money is 10^{2^n} . Sorry Bernoulli! **QED**

There are *a lot* of ornate but unsuccessful proposed solutions.

The St. Petersburg Paradox: A Subjective Probability Solution

Hongbin Wang (hwang@tamhsc.edu)

Yanlong Sun (ysun@tamhsc.edu)

Jack W. Smith (jwsmith@tamhsc.edu)

Texas A&M University Health Science Center

2121 West Holcombe Blvd, Suite 1109, Houston, TX 77030 USA

Abstract

The St. Petersburg Paradox (SPP), where people are willing to pay only a modest amount for a lottery with infinite expected gain, has been a famous showcase of human (ir)rationality. Since inception multiple solutions have been proposed, including the influential expected utility theory. Criticisms remain due to the lack of a priori justification for the utility function. Here we report a new solution to the long-standing paradox, which focuses on the probability weighting component (rather than the value/utility component) in calculating the expected value of the game. We show that a new Additional Transition Time (AT) based measure, motivated by both physics and psychology, can naturally lead to a converging expected value and therefore solve the paradox.

Keywords: human judgment and decision making, probability, St. Petersburg Paradox,

Fate laughs at probabilities.

-- E. G. Bulwer-Lytton

Introduction

Suppose you are offered the following gamble:

- Toss a fair coin. If you get a head, you are paid \$1 and the game is over. Otherwise, toss again.
- If you get a head in the second tossing, you are paid \$2 and the game is over. Otherwise, toss again.
- If you get a head in the third tossing, you are paid \$4 and the game is over. Otherwise, toss again.
- ... Game continues until you get a head. If you get a head in the n th tossing, you will be paid $\$2^{n-1}$.

How much are you willing to pay to play this gamble?

A simple calculation shows that the gamble's expected value, S , is infinite:

$$S = \$ \sum_{n=1}^{\infty} p^n 2^{n-1} = \$ \sum_{n=1}^{\infty} \left(\frac{1}{2}\right)^n 2^{n-1} = \$ \left(\frac{1}{2} + \frac{1}{2} + \dots\right) \quad \text{Eq. 1}$$

where n is the number of tosses to get the first head (i.e., after a streak of $n-1$ tails, one gets a head, and the game is over).

The question is, are you willing to pay any price for a right to play this game? Probably not. More than three hundred years ago, in 1713, Nicolas Bernoulli, a young Swiss mathematician, first proposed this problem and pointed out that a sensible person would only be willing to pay very little to play the game. This constitutes a contradiction, which nowadays is called the St. Petersburg Paradox (SPP).

A Little History

The SPP was so named after the eponymous Russian city, where Daniel Bernoulli, a mathematician and Nicholas Bernoulli's cousin, published his classical solution to the problem in 1738. However, the problem was initially proposed by Nicolas Bernoulli in 1713, who was clearly troubled by it. According to him, while the expectation of game gain was infinity, the player would be guaranteed to lose since it is "morally impossible" that one not achieve a head in a finite number of tossing.

In 1728, Gabriel Cramer, another Swiss mathematician, wrote to N. Bernoulli and suggested a solution. In Cramer's solution, money's quantity was replaced by its "moral value", representing the pleasure or sorrow money (or loss of money) could produce. In doing so Cramer showed the expectation would converge to less than \$3 if "one wishes to suppose that the moral value of goods was as the square root of the mathematical quantities".

N. Bernoulli was not entirely satisfied with this solution. In his reply to Cramer, N. Bernoulli wrote that the pleasure difference "does not demonstrate the true reason" for why one should not pay infinity to play the game. Even Cramer himself thought his square-root assumption about money and pleasure was not just.

Eventually in 1738, D. Bernoulli published his solution to the problem (Bernoulli, 1738). D. Bernoulli's solution was similar to Cramer's and based on the concept of utility, which measured the usefulness of values and was taken to be a logarithmic function of values. It was shown that while the expected value diverged the expected utility converged. D. Bernoulli's solution was seminal and extremely influential, and has since shaped the whole field of economics and of the psychology of decision making.

It was interesting to note that N. Bernoulli vigorously objected his cousin's approach. A series of communication showed that the two had engaged in serious arguments. To N. Bernoulli, the concept of utility, similar to the "moral value" of Cramer, was arbitrary and, to a certain extent, irrelevant. Rather, the concern here was to find a more general way to show if a game was fair, regardless of who was playing the game. "For example a game is considered fair, when the two players bet an equal sum on a game under equal conditions, although according to your theory, and by paying attention to their riches, the pleasure or the advantage of gain in the favorable case is not equal to the

Solution

Solution

- I. It's irrational to bank on things that are *beyond reasonable belief*; you know this.

Solution

1. It's irrational to bank on things that are *beyond reasonable belief*; you know this.
2. It's *beyond reasonable belief* that you'll win back our \$30 price to play; you know this too.

Solution

1. It's irrational to bank on things that are *beyond reasonable belief*; you know this.
2. It's *beyond reasonable belief* that you'll win back our \$30 price to play; you know this too.
3. Hence, since you are rational, you'll decline.

Solution

1. It's irrational to bank on things that are *beyond reasonable belief*; you know this.
2. It's *beyond reasonable belief* that you'll win back our \$30 price to play; you know this too.
3. Hence, since you're rational, you'll decline.

Solution

1. It's irrational to bank on things that are *beyond reasonable belief*; you know this.
2. It's *beyond reasonable belief* that you'll win back our \$30 price to play; you know this too.
3. Hence, since you're rational, you'll decline.

It's *beyond reasonable belief* that \$32 will be won (an amount that would make a bet of \$30 a nice wager), since the odds of that happening are $1/32$.

Solution

1. It's irrational to bank on things that are *beyond reasonable belief*; you know this.
2. It's *beyond reasonable belief* that you'll win back our \$30 price to play; you know this too.
3. Hence, since you're rational, you'll decline.

It's *beyond reasonable belief* that \$32 will be won (an amount that would make a bet of \$30 a nice wager), since the odds of that happening are $1/32$.

Symmetrically put, a rational agent will believe at level 3 that \$30 dollars will be lost. (Why?)

Solution

1. It's irrational to bank on things that are *beyond reasonable belief*; you know this.
2. It's *beyond reasonable belief* that you'll win back our \$30 price to play; you know this too.
3. Hence, since you're rational, you'll decline.

It's *beyond reasonable belief* that \$32 will be won (an amount that would make a bet of \$30 a nice wager), since the odds of that happening are $1/32$.

Symmetrically put, a rational agent will believe at level 3 that \$30 dollars will be lost. (Why?)

The needed principle: *Ceteris paribus*, if a rational agent *a* believes at level 3 that a wager cutting meaningfully into that agent's disposable income will be lost, *a* will not make that wager.

Solution

1. It's irrational to bank on things that are *beyond reasonable belief*; you know this.
2. It's *beyond reasonable belief* that you'll win back our \$30 price to play; you know this too.
3. Hence, since you're rational, you'll decline.

It's *beyond reasonable belief* that \$32 will be won (an amount that would make a bet of \$30 a nice wager), since the odds of that happening are 1/32.

Symmetrically put, a rational agent will believe at level 3 that \$30 dollars will be lost. (Why?)

The needed principle: *Ceteris paribus*, if a rational agent *a* believes at level 3 that a wager cutting meaningfully into that agent's disposable income will be lost, *a* will not make that wager.

From SEP entry: "... few of us would pay even \$25 to enter such a game." Now we know why!

Paradoxes are engines of progress in formal logic.

E.g., Russell's Paradox — as we've seen.

The paradox in this case?

The paradox in this case?

- A perfectly rational person can never believe P and believe $\neg P$ at the same time.

The paradox in this case?

- A perfectly rational person can never believe P and believe $\neg P$ at the same time.
- The Lottery Paradox (apparently) shows, courtesy of its two Sequences (of Reasoning), that a perfectly rational person can indeed have such a belief (upon considering a fair, large lottery).

The paradox in this case?

- A perfectly rational person can never believe P and believe $\neg P$ at the same time.
- The Lottery Paradox (apparently) shows, courtesy of its two Sequences (of Reasoning), that a perfectly rational person can indeed have such a belief (upon considering a fair, large lottery).
- Contradiction! — and hence a paradox!

Types of Paradoxes

- Deductive Paradoxes. The reasoning in question is exclusively deductive.
 - Russell's Paradox
 - The Liar Paradox
 - Richard's Paradox
- Inductive Paradoxes. Some of the reasoning in question uses non-deductive reasoning (e.g., probabilistic reasoning, abductive reasoning, analogical reasoning, etc.).

Types of Paradoxes

- Deductive Paradoxes. The reasoning in question is exclusively deductive.
 - Russell's Paradox
 - The Liar Paradox
 - Richard's Paradox
- Inductive Paradoxes Some of the reasoning in question uses non-deductive reasoning (e.g., probabilistic reasoning, abductive reasoning, analogical reasoning, etc.).

Types of Paradoxes

- Deductive Paradoxes. The reasoning in question is exclusively deductive.
 - Russell's Paradox
 - The Liar Paradox
 - Richard's Paradox
- Inductive Paradoxes. Some of the reasoning in question uses non-deductive reasoning (e.g., probabilistic reasoning, abductive reasoning, analogical reasoning, etc.).

Types of Paradoxes

- Deductive Paradoxes. The reasoning in question is exclusively deductive.
 - Russell's Paradox
 - The Liar Paradox
 - Richard's Paradox
- Inductive Paradoxes Some of the reasoning in question uses non-deductive reasoning (e.g., probabilistic reasoning, abductive reasoning, analogical reasoning, etc.).

Inductive Logic

the hallmark of deductive logic is *proof*, the hallmark of inductive logic is the concept of an *argument*. An exceptionally strong kind of argument is a proof, but plenty of arguments fall short of being proofs — and yet still have considerable force. For instance, consider the following argument α_1 :

- (1) Tweety is bird.
 - (2) Most birds can fly.
-
- \therefore (3) Tweety can fly.

For stark contrast, consider as well this argument (α_2):

- (1') 3 is a positive integer.
 - (2') All positive integers are greater than 0.
-
- \therefore (3') 3 is greater than 0.

The second of these arguments qualifies as an outright proof. That is, using the notation much employed before the present chapter:

$$\{(1'), (2')\} \vdash (3')$$

But in stark contrast, argument α_1 is not a proof that Tweety can fly. The reason is obvious: (3) isn't deduced from the combination of (1) and (2); that is,

$$\{(1), (2)\} \not\vdash (3)$$

Inductive Logic

the hallmark of deductive logic is *proof*, the hallmark of inductive logic is the concept of an *argument*. An exceptionally strong kind of argument is a proof, but plenty of arguments fall short of being proofs — and yet still have considerable force. For instance, consider the following argument α_1 :

- (1) Tweety is bird.
- (2) Most birds can fly.

- \therefore (3) Tweety can fly.

For stark contrast, consider as well this argument (α_2):

- (1') 3 is a positive integer.
- (2') All positive integers are greater than 0.

- \therefore (3') 3 is greater than 0.

The second of these arguments qualifies as an outright proof. That is, using the notation much employed before the present chapter:

$$\{(1'), (2')\} \vdash (3')$$

But in stark contrast, argument α_1 is not a proof that Tweety can fly. The reason is obvious: (3) isn't deduced from the combination of (1) and (2); that is,

$$\{(1), (2)\} \not\vdash (3)$$

Inductive Logic

the hallmark of deductive logic is *proof*, the hallmark of inductive logic is the concept of an *argument*. An exceptionally strong kind of argument is a proof, but plenty of arguments fall short of being proofs — and yet still have considerable force. For instance, consider the following argument α_1 :

- | | |
|------------------|---------------------|
| (1) | Tweety is bird. |
| (2) | Most birds can fly. |
| <hr/> | |
| \therefore (3) | Tweety can fly. |

For stark contrast, consider as well this argument (α_2):

- | | |
|-------------------|---|
| (1') | 3 is a positive integer. |
| (2') | All positive integers are greater than 0. |
| <hr/> | |
| \therefore (3') | 3 is greater than 0. |

Deductive (familiar)

The second of these arguments qualifies as an outright proof. That is, using the notation much employed before the present chapter:

$$\{(1'), (2')\} \vdash (3')$$

But in stark contrast, argument α_1 is not a proof that Tweety can fly. The reason is obvious: (3) isn't deduced from the combination of (1) and (2); that is,

$$\{(1), (2)\} \not\vdash (3)$$

Inductive Logic

the hallmark of deductive logic is *proof*, the hallmark of inductive logic is the concept of an *argument*. An exceptionally strong kind of argument is a proof, but plenty of arguments fall short of being proofs — and yet still have considerable force. For instance, consider the following argument α_1 :

(1)	Tweety is bird.
(2)	Most birds can fly.
<hr/>	
\therefore (3)	Tweety can fly.

For stark contrast, consider as well this argument (α_2):

(1')	3 is a positive integer.
(2')	All positive integers are greater than 0.
<hr/>	
\therefore (3')	3 is greater than 0.

Deductive (familiar)

The second of these arguments qualifies as an outright proof. That is, using the notation much employed before the present chapter:

$$\{(1'), (2')\} \vdash (3')$$

But in stark contrast, argument α_1 is not a proof that Tweety can fly. The reason is obvious: (3) isn't deduced from the combination of (1) and (2); that is,

$$\{(1), (2)\} \not\vdash (3)$$

Inductive Logic

the hallmark of deductive logic is *proof*, the hallmark of inductive logic is the concept of an *argument*. An exceptionally strong kind of argument is a proof, but plenty of arguments fall short of being proofs — and yet still have considerable force. For instance, consider the following argument α_1 :

(1)	Tweety is bird.
(2)	Most birds can fly.
<hr/>	
\therefore (3)	Tweety can fly.

Inductive (new)

For stark contrast, consider as well this argument (α_2):

(1')	3 is a positive integer.
(2')	All positive integers are greater than 0.
<hr/>	
\therefore (3')	3 is greater than 0.

Deductive (familiar)

The second of these arguments qualifies as an outright proof. That is, using the notation much employed before the present chapter:

$$\{(1'), (2')\} \vdash (3')$$

But in stark contrast, argument α_1 is not a proof that Tweety can fly. The reason is obvious: (3) isn't deduced from the combination of (1) and (2); that is,

$$\{(1), (2)\} \not\vdash (3)$$

Inductive-Reasoning Example from Pollock — for Peek Ahead

Imagine the following:

Keith tells you that the morning news predicts rain in Troy today. However, Alvin tells you that the same news report predicted sunshine.

Imagine the following: Keith tells you that the morning news predicts rain in Tucson today. However, Alvin tells you that the same news report predicted sunshine.

Without any other source of information, it would be irrational to place belief in either Keith's or Alvin's statements.

Imagine the following: Keith tells you that the morning news predicts rain in Tucson today. However, Alvin tells you that the same news report predicted sunshine.

Without any other source of information, it would be irrational to place belief in either Keith's or Alvin's statements.

Further, suppose you happened to watch the noon news report, and that report predicted rain. Then you should believe that it will rain despite your knowledge of Alvin's argument.

Defeasible Reasoning in OSCAR


K- Keith says that M

A- Alvin says that $\sim M$

M- The morning news said that R

R- It is going to rain this afternoon

N- The noon news says that R



All such can be absorbed into our inductive logics and our automated inductive reasoners (= our AI).

In Our Inductive Modal Logic

- | | | |
|-----|--|------|
| (1) | $\mathbf{K}(you, \mathbf{S}(keith, \mathbf{S}(m, rain)))$ | fact |
| (2) | $\mathbf{K}(you, \mathbf{S}(alvin, \mathbf{S}(m, \neg rain)))$ | fact |

In Our Inductive Modal Logic

	(1)	$\mathbf{K}(you, \mathbf{S}(keith, \mathbf{S}(m, rain)))$	fact
	(2)	$\mathbf{K}(you, \mathbf{S}(alvin, \mathbf{S}(m, \neg rain)))$	fact
\therefore	(3)	$\mathbf{S}(keith, \mathbf{S}(m, rain))$?
\therefore	(4)	$\mathbf{S}(alvin, \mathbf{S}(m, \neg rain))$?

In Our Inductive Modal Logic

	(1)	$\mathbf{K}(you, \mathbf{S}(keith, \mathbf{S}(m, rain)))$	fact
	(2)	$\mathbf{K}(you, \mathbf{S}(alvin, \mathbf{S}(m, \neg rain)))$	fact
\therefore	(3)	$\mathbf{S}(keith, \mathbf{S}(m, rain))$?
\therefore	(4)	$\mathbf{S}(alvin, \mathbf{S}(m, \neg rain))$?
	(5)	$\mathbf{S}(keith, \phi) \rightarrow \mathbf{B}^2(you, \phi)$	Testimonial P1

In Our Inductive Modal Logic

	(1)	$\mathbf{K}(you, \mathbf{S}(keith, \mathbf{S}(m, rain)))$	fact
	(2)	$\mathbf{K}(you, \mathbf{S}(alvin, \mathbf{S}(m, \neg rain)))$	fact
\therefore	(3)	$\mathbf{S}(keith, \mathbf{S}(m, rain))$?
\therefore	(4)	$\mathbf{S}(alvin, \mathbf{S}(m, \neg rain))$?
	(5)	$\mathbf{S}(keith, \phi) \rightarrow \mathbf{B}^2(you, \phi)$	Testimonial P1
\therefore	(6)	$\mathbf{B}^2(you, \mathbf{S}(m, rain)) \wedge \mathbf{B}^2(you, \mathbf{S}(m, \neg rain))$	

In Our Inductive Modal Logic

	(1)	$\mathbf{K}(you, \mathbf{S}(keith, \mathbf{S}(m, rain)))$	fact
	(2)	$\mathbf{K}(you, \mathbf{S}(alvin, \mathbf{S}(m, \neg rain)))$	fact
\therefore	(3)	$\mathbf{S}(keith, \mathbf{S}(m, rain))$?
\therefore	(4)	$\mathbf{S}(alvin, \mathbf{S}(m, \neg rain))$?
	(5)	$\mathbf{S}(keith, \phi) \rightarrow \mathbf{B}^2(you, \phi)$	Testimonial P1
\therefore	(6)	$\mathbf{B}^2(you, \mathbf{S}(m, rain)) \wedge \mathbf{B}^2(you, \mathbf{S}(m, \neg rain))$	
\therefore	(7)	$\neg \mathbf{B}^2(you, \mathbf{S}(m, rain)) \wedge \neg \mathbf{B}^2(you, \mathbf{S}(m, \neg rain))$	“Clash” Principle

In Our Inductive Modal Logic

(1)	$\mathbf{K}(you, \mathbf{S}(keith, \mathbf{S}(m, rain)))$	fact
(2)	$\mathbf{K}(you, \mathbf{S}(alvin, \mathbf{S}(m, \neg rain)))$	fact
\therefore	(3) $\mathbf{S}(keith, \mathbf{S}(m, rain))$?
\therefore	(4) $\mathbf{S}(alvin, \mathbf{S}(m, \neg rain))$?
(5)	$\mathbf{S}(keith, \phi) \rightarrow \mathbf{B}^2(you, \phi)$	Testimonial P1
\therefore	(6) $\mathbf{B}^2(you, \mathbf{S}(m, rain)) \wedge \mathbf{B}^2(you, \mathbf{S}(m, \neg rain))$	
\therefore	(7) $\neg \mathbf{B}^2(you, \mathbf{S}(m, rain)) \wedge \neg \mathbf{B}^2(you, \mathbf{S}(m, \neg rain))$	“Clash” Principle
(8)	$\mathbf{K}(you, \mathbf{S}(noonnews, rain))$	

In Our Inductive Modal Logic

(1)	$\mathbf{K}(you, \mathbf{S}(keith, \mathbf{S}(m, rain)))$	fact
(2)	$\mathbf{K}(you, \mathbf{S}(alvin, \mathbf{S}(m, \neg rain)))$	fact
\therefore	(3) $\mathbf{S}(keith, \mathbf{S}(m, rain))$?
\therefore	(4) $\mathbf{S}(alvin, \mathbf{S}(m, \neg rain))$?
(5)	$\mathbf{S}(keith, \phi) \rightarrow \mathbf{B}^2(you, \phi)$	Testimonial P1
\therefore	(6) $\mathbf{B}^2(you, \mathbf{S}(m, rain)) \wedge \mathbf{B}^2(you, \mathbf{S}(m, \neg rain))$	
\therefore	(7) $\neg \mathbf{B}^2(you, \mathbf{S}(m, rain)) \wedge \neg \mathbf{B}^2(you, \mathbf{S}(m, \neg rain))$	“Clash” Principle
(8)	$\mathbf{K}(you, \mathbf{S}(noonnews, rain))$	
\therefore	(9) $\mathbf{S}(noonnews, rain)$?

In Our Inductive Modal Logic

	(1)	$\mathbf{K}(you, \mathbf{S}(keith, \mathbf{S}(m, rain)))$	fact
	(2)	$\mathbf{K}(you, \mathbf{S}(alvin, \mathbf{S}(m, \neg rain)))$	fact
\therefore	(3)	$\mathbf{S}(keith, \mathbf{S}(m, rain))$?
\therefore	(4)	$\mathbf{S}(alvin, \mathbf{S}(m, \neg rain))$?
	(5)	$\mathbf{S}(keith, \phi) \rightarrow \mathbf{B}^2(you, \phi)$	Testimonial P1
\therefore	(6)	$\mathbf{B}^2(you, \mathbf{S}(m, rain)) \wedge \mathbf{B}^2(you, \mathbf{S}(m, \neg rain))$	
\therefore	(7)	$\neg \mathbf{B}^2(you, \mathbf{S}(m, rain)) \wedge \neg \mathbf{B}^2(you, \mathbf{S}(m, \neg rain))$	“Clash” Principle
	(8)	$\mathbf{K}(you, \mathbf{S}(noonnews, rain))$	
\therefore	(9)	$\mathbf{S}(noonnews, rain)$?
	(10)	$\mathbf{S}(noonnews, \phi) \rightarrow \mathbf{B}^3(you, \phi)$	Testimonial P2

In Our Inductive Modal Logic

(1)	$\mathbf{K}(you, \mathbf{S}(keith, \mathbf{S}(m, rain)))$	fact
(2)	$\mathbf{K}(you, \mathbf{S}(alvin, \mathbf{S}(m, \neg rain)))$	fact
\therefore	(3) $\mathbf{S}(keith, \mathbf{S}(m, rain))$?
\therefore	(4) $\mathbf{S}(alvin, \mathbf{S}(m, \neg rain))$?
(5)	$\mathbf{S}(keith, \phi) \rightarrow \mathbf{B}^2(you, \phi)$	Testimonial P1
\therefore	(6) $\mathbf{B}^2(you, \mathbf{S}(m, rain)) \wedge \mathbf{B}^2(you, \mathbf{S}(m, \neg rain))$	
\therefore	(7) $\neg \mathbf{B}^2(you, \mathbf{S}(m, rain)) \wedge \neg \mathbf{B}^2(you, \mathbf{S}(m, \neg rain))$	“Clash” Principle
(8)	$\mathbf{K}(you, \mathbf{S}(noonnews, rain))$	
\therefore	(9) $\mathbf{S}(noonnews, rain)$?
(10)	$\mathbf{S}(noonnews, \phi) \rightarrow \mathbf{B}^3(you, \phi)$	Testimonial P2
\therefore	(11) $\mathbf{B}^3(you, rain)$	

The Lottery Paradox ...





E: “Please go down to Stewart’s & get the T U.”



E: “Please go down to Stewart’s & get the T U.”







S: “I’m sorry, E, I’m afraid I can’t do that.
It would be irrational.”











E: “Please go down to Stewart’s & get the T U.”

S: “I’m sorry, E, I’m afraid I can’t do that.
It would be irrational.”


•••




Sequence I




Sequence I




Sequence I




Sequence I




Sequence I




Sequence I







Sequence 1




Sequence 2




Sequence 1




Sequence 2




Sequence 1




Sequence 2





Sequence 1



Sequence 2







Sequence 1

Sequence 2

Contradiction!

Sequence I

Sequence I

Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent a is fully apprised.

Sequence I

Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

Sequence I

Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

$$Wt_1 \oplus Wt_2 \oplus \dots \oplus Wt_{1T} \quad (1)$$

Sequence I

Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

$$Wt_1 \oplus Wt_2 \oplus \dots \oplus Wt_{1T} \quad (1)$$

We then deduce from this that there is at least one ticket that will win, a proposition represented — using standard notation — as:

Sequence I

Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

$$Wt_1 \oplus Wt_2 \oplus \dots \oplus Wt_{1T} \quad (1)$$

We then deduce from this that there is at least one ticket that will win, a proposition represented — using standard notation — as:

$$\exists t_i Wt_i \quad (2)$$

Sequence I

Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

$$Wt_1 \oplus Wt_2 \oplus \dots \oplus Wt_{1T} \quad (1)$$

We then deduce from this that there is at least one ticket that will win, a proposition represented — using standard notation — as:

$$\exists t_i Wt_i \quad (2)$$

Very well; perfectly clear so far. And now we can add another deductive step: Since our rational agent *a* can follow this deduction sequence to this point, and since **D** is assumed to be indubitable, it follows that our rational agent in turn believes (2); i.e., we conclude Sequence I by obtaining the following:

Sequence I

Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

$$Wt_1 \oplus Wt_2 \oplus \dots \oplus Wt_{1T} \quad (1)$$


We then deduce from this that there is at least one ticket that will win, a proposition represented — using standard notation — as:

$$\exists t_i Wt_i \quad (2)$$


Very well; perfectly clear so far. And now we can add another deductive step: Since our rational agent *a* can follow this deduction sequence to this point, and since **D** is assumed to be indubitable, it follows that our rational agent in turn believes (2); i.e., we conclude Sequence I by obtaining the following:

$$\mathbf{B}_a \exists t_i Wt_i \quad (3)$$

Sequence I



Sequence I



Sequence 2

Sequence 2

As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

Sequence 2

As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using 'IT' to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

Sequence 2

As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using '1T' to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$\text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

Sequence 2

As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$\text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_i winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet. Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_i won't win sails through—and this of course works for each ticket. Hence we have:

Sequence 2

As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$\text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet. Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:

$$\mathbf{B}_a \neg Wt_1 \wedge \mathbf{B}_a \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a \neg Wt_{1T} \quad (2)$$

Sequence 2

As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using '1T' to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$\text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet. Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:

$$\mathbf{B}_a \neg Wt_1 \wedge \mathbf{B}_a \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a \neg Wt_{1T} \quad (2)$$

Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:

Sequence 2

As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using '1T' to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$\text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_i winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet. Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_i won't win sails through—and this of course works for each ticket. Hence we have:

$$\mathbf{B}_a \neg Wt_1 \wedge \mathbf{B}_a \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a \neg Wt_{1T} \quad (2)$$

Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:

$$\mathbf{B}_a (\neg Wt_1 \wedge \neg Wt_2 \wedge \dots \wedge \neg Wt_{1T}) \quad (3)$$

Sequence 2

As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$\text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet. Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:

$$\mathbf{B}_a \neg Wt_1 \wedge \mathbf{B}_a \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a \neg Wt_{1T} \quad (2)$$

Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:

$$\mathbf{B}_a (\neg Wt_1 \wedge \neg Wt_2 \wedge \dots \wedge \neg Wt_{1T}) \quad (3)$$

But (3) is logically equivalent to the statement that there doesn't exist a winning ticket; i.e., (3) is logically equivalent to this result from Sequence 2:

Sequence 2

As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using '1T' to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$\text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet. Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:


$$\mathbf{B}_a \neg Wt_1 \wedge \mathbf{B}_a \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a \neg Wt_{1T} \quad (2)$$


Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:


$$\mathbf{B}_a (\neg Wt_1 \wedge \neg Wt_2 \wedge \dots \wedge \neg Wt_{1T}) \quad (3)$$


But (3) is logically equivalent to the statement that there doesn't exist a winning ticket; i.e., (3) is logically equivalent to this result from Sequence 2:


$$\mathbf{B}_a \neg \exists t_i Wt_i \quad (4)$$











(The contradiction we sketched earlier has arrived.)

A Solution to The Lottery Paradox ...

Strength-Factor Continuum

Certain

Improbable

Evidently False

Probable

Beyond Reasonable Belief

Certainly False

Counterbalanced

Evident

Beyond Reasonable Doubt

Strength-Factor Continuum

Certain

Evident

Beyond Reasonable Doubt

Probable

Counterbalanced

Improbable

Beyond Reasonable Belief

Evidently False

Certainly False

Actually, now ...

Actually, now ...

English	Value
certain	6
evident	5
overwhelmingly likely = “beyond reasonable doubt” = “one in a million”	4
very likely	3
likely	2
more likely than not	1
counterbalanced	0

Actually, now ...

English	Value
certain	6
evident	5
overwhelmingly likely = “beyond reasonable doubt” = “one in a million”	4
very likely	3
likely	2
more likely than not	1
counterbalanced	0

... but let's use the simpler scheme.

Strength-Factor Continuum

Certain

Evident

Beyond Reasonable Doubt

Probable

Counterbalanced

Improbable

Beyond Reasonable Belief

Evidently False

Certainly False

Strength-Factor Continuum

Certain

Evident

Beyond Reasonable Doubt

Probable

..... Counterbalanced

Improbable

Beyond Reasonable Belief

Evidently False

Certainly False

Strength-Factor Continuum

Epistemically Positive

Certain

Evident

Beyond Reasonable Doubt

Probable

..... Counterbalanced

Improbable

Beyond Reasonable Belief

Evidently False

Certainly False

Strength-Factor Continuum

Epistemically Positive

Certain

Evident

Beyond Reasonable Doubt

Probable

..... Counterbalanced

Improbable

Beyond Reasonable Belief

Evidently False

Epistemically Negative

Certainly False

Strength-Factor Continuum

Epistemically Positive

Certain

Evident

Beyond Reasonable Doubt

Probable

.....

Counterbalanced

Improbable

Beyond Reasonable Belief

Evidently False

Epistemically Negative

Certainly False

Strength-Factor Continuum

Epistemically Positive

(4) Certain

(3) Evident

(2) Beyond Reasonable Doubt

(1) Probable

..... (0) Counterbalanced

(-1) Improbable

(-2) Beyond Reasonable Belief

(-3) Evidently False

Epistemically Negative

(-4) Certainly False

Strength-Factor Continuum

Epistemically Positive

(4) Certain

(3) Evident

(2) Beyond Reasonable Doubt

(1) Probable

..... (0) Counterbalanced

(-1) Improbable

(-2) Beyond Reasonable Belief

(-3) Evidently False

Epistemically Negative

(-4) Certainly False

Epistemically Positive

(4) Certain

(3) Evident

(2) Beyond Reasonable Doubt

(1) Probable

..... (0) Counterbalanced

(-1) Improbable

(-2) Beyond Reasonable Belief

(-3) Evidently False

Epistemically Negative

(-4) Certainly False

Key Principles

Epistemically Positive

(4) Certain

(3) Evident

(2) Beyond Reasonable Doubt

(1) Probable

..... (0) Counterbalanced

(-1) Improbable

(-2) Beyond Reasonable Belief

(-3) Evidently False

Epistemically Negative

(-4) Certainly False

Key Principles

Epistemically Positive

Deduction preserves strength.

(4) Certain

(3) Evident

(2) Beyond Reasonable Doubt

(1) Probable

..... (0) Counterbalanced

(-1) Improbable

(-2) Beyond Reasonable Belief

(-3) Evidently False

Epistemically Negative

(-4) Certainly False

Key Principles

Epistemically Positive

(4) Certain

Deduction preserves strength.

(3) Evident

Clashes are resolved in favor of higher strength.

(2) Beyond Reasonable Doubt

(1) Probable

..... (0) Counterbalanced

(-1) Improbable

(-2) Beyond Reasonable Belief

(-3) Evidently False

Epistemically Negative

(-4) Certainly False

Key Principles

Epistemically Positive

(4) Certain

Deduction preserves strength.

(3) Evident

Clashes are resolved in favor of higher strength.

(2) Beyond Reasonable Doubt

(1) Probable

..... **Any proposition p such that $\text{prob}(p) < 1$ is at most evident.**

(0) Counterbalanced

(-1) Improbable

(-2) Beyond Reasonable Belief

(-3) Evidently False

Epistemically Negative

(-4) Certainly False

Key Principles

Epistemically Positive

(4) Certain

Deduction preserves strength.

(3) Evident

Clashes are resolved in favor of higher strength.

(2) Beyond Reasonable Doubt

(1) Probable

..... **Any proposition p such that $\text{prob}(p) < 1$ is at most evident.**

(0) Credible

Any rational belief that p , where the basis for p is at most evident, is at most an evident (= level 3) belief.

(-1) Improbable

(-2) Beyond Reasonable Doubt Belief

(-3) Evidently False

Epistemically Negative

(-4) Certainly False

Sequence I, “Rigorized”

Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

$$Wt_1 \oplus Wt_2 \oplus \dots \oplus Wt_{1T} \quad (1)$$

We then deduce from this that there is at least one ticket that will win, a proposition represented — using standard notation — as:

$$\exists t_i Wt_i \quad (2)$$

Very well; perfectly clear so far. And now we can add another deductive step: Since our rational agent *a* can follow this deduction sequence to this point, and since **D** is assumed to be indubitable, it follows that our rational agent in turn believes (2); i.e., we conclude Sequence I by obtaining the following:

$$\mathbf{B}_a \exists t_i Wt_i \quad (3)$$

Sequence I, “Rigorized”

- 4 Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

$$Wt_1 \oplus Wt_2 \oplus \dots \oplus Wt_{1T} \quad (1)$$

We then deduce from this that there is at least one ticket that will win, a proposition represented — using standard notation — as:

$$\exists t_i Wt_i \quad (2)$$

Very well; perfectly clear so far. And now we can add another deductive step: Since our rational agent *a* can follow this deduction sequence to this point, and since **D** is assumed to be indubitable, it follows that our rational agent in turn believes (2); i.e., we conclude Sequence I by obtaining the following:

$$\mathbf{B}_a \exists t_i Wt_i \quad (3)$$

Sequence I, “Rigorized”

- 4 Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

$$4 \quad Wt_1 \oplus Wt_2 \oplus \dots \oplus Wt_{1T} \quad (1)$$

We then deduce from this that there is at least one ticket that will win, a proposition represented — using standard notation — as:

$$\exists t_i Wt_i \quad (2)$$

Very well; perfectly clear so far. And now we can add another deductive step: Since our rational agent *a* can follow this deduction sequence to this point, and since **D** is assumed to be indubitable, it follows that our rational agent in turn believes (2); i.e., we conclude Sequence I by obtaining the following:

$$\mathbf{B}_a \exists t_i Wt_i \quad (3)$$

Sequence I, “Rigorized”

- 4 Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

$$4 \quad Wt_1 \oplus Wt_2 \oplus \dots \oplus Wt_{1T} \quad (1)$$

We then deduce from this that there is at least one ticket that will win, a proposition represented — using standard notation — as:

$$4 \quad \exists t_i Wt_i \quad (2)$$

Very well; perfectly clear so far. And now we can add another deductive step: Since our rational agent *a* can follow this deduction sequence to this point, and since **D** is assumed to be indubitable, it follows that our rational agent in turn believes (2); i.e., we conclude Sequence I by obtaining the following:

$$B_a \exists t_i Wt_i \quad (3)$$

Sequence I, “Rigorized”

- 4 Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

$$4 \quad Wt_1 \oplus Wt_2 \oplus \dots \oplus Wt_{1T} \quad (1)$$

We then deduce from this that there is at least one ticket that will win, a proposition represented — using standard notation — as:

$$4 \quad \exists t_i Wt_i \quad (2)$$

Very well; perfectly clear so far. And now we can add another deductive step: Since our rational agent *a* can follow this deduction sequence to this point, and since **D** is assumed to be indubitable, it follows that our rational agent in turn believes (2); i.e., we conclude Sequence I by obtaining the following:

$$4 \quad B_a \exists t_i Wt_i \quad (3)$$

Sequence I, “Rigorized”

- 4 Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

$$4 \quad Wt_1 \oplus Wt_2 \oplus \dots \oplus Wt_{1T} \quad (1)$$

We then deduce from this that there is at least one ticket that will win, a proposition represented — using standard notation — as:

$$4 \quad \exists t_i Wt_i \quad (2)$$

Very well; perfectly clear so far. And now we can add another deductive step: Since our rational agent *a* can follow this deduction sequence to this point, and since **D** is assumed to be indubitable, it follows that our rational agent in turn believes (2); i.e., we conclude Sequence I by obtaining the following:

Sequence I, “Rigorized”

- 4 Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

$$4 \quad Wt_1 \oplus Wt_2 \oplus \dots \oplus Wt_{1T} \quad (1)$$

We then deduce from this that there is at least one ticket that will win, a proposition represented — using standard notation — as:

$$4 \quad \exists t_i Wt_i \quad (2)$$

Very well; perfectly clear so far. And now we can add another deductive step: Since our rational agent *a* can follow this deduction sequence to this point, and since **D** is assumed to be indubitable, it follows that our rational agent in turn believes (2); i.e., we conclude Sequence I by obtaining the following:

$$4 \quad B_a^4 \exists t_i Wt_i \quad (3)$$

Sequence 2, “Rigorized”

As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$\text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet.

Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:

$$\mathbf{B}_a \neg Wt_1 \wedge \mathbf{B}_a \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a \neg Wt_{1T} \quad (2)$$

Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:

$$\mathbf{B}_a (\neg Wt_1 \wedge \neg Wt_2 \wedge \dots \wedge \neg Wt_{1T}) \quad (3)$$

But (3) is logically equivalent to the statement that there doesn't exist a winning ticket; i.e., (3) is logically equivalent to this result from Sequence 2:

$$\mathbf{B}_a \neg \exists t_i Wt_i \quad (4)$$

Sequence 2, “Rigorized”

4

As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$\text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet.

Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:

$$\mathbf{B}_a \neg Wt_1 \wedge \mathbf{B}_a \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a \neg Wt_{1T} \quad (2)$$

Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:

$$\mathbf{B}_a (\neg Wt_1 \wedge \neg Wt_2 \wedge \dots \wedge \neg Wt_{1T}) \quad (3)$$

But (3) is logically equivalent to the statement that there doesn't exist a winning ticket; i.e., (3) is logically equivalent to this result from Sequence 2:

$$\mathbf{B}_a \neg \exists t_i Wt_i \quad (4)$$

Sequence 2, “Rigorized”

4 As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$4 \quad \text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet. Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:

$$\mathbf{B}_a \neg Wt_1 \wedge \mathbf{B}_a \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a \neg Wt_{1T} \quad (2)$$

Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:

$$\mathbf{B}_a (\neg Wt_1 \wedge \neg Wt_2 \wedge \dots \wedge \neg Wt_{1T}) \quad (3)$$

But (3) is logically equivalent to the statement that there doesn't exist a winning ticket; i.e., (3) is logically equivalent to this result from Sequence 2:

$$\mathbf{B}_a \neg \exists t_i Wt_i \quad (4)$$

Sequence 2, “Rigorized”

4 As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$4 \quad \text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet. Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:

Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:

$$\mathbf{B}_a(\neg Wt_1 \wedge \neg Wt_2 \wedge \dots \wedge \neg Wt_{1T}) \quad (3)$$

But (3) is logically equivalent to the statement that there doesn't exist a winning ticket; i.e., (3) is logically equivalent to this result from Sequence 2:

$$\mathbf{B}_a \neg \exists t_i Wt_i \quad (4)$$

Sequence 2, “Rigorized”

4 As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$4 \quad \text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet.

Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:

$$\mathbf{B}_a^3 \neg Wt_1 \wedge \mathbf{B}_a^3 \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a^3 \neg Wt_{1T} \quad (2)$$

Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:

$$\mathbf{B}_a(\neg Wt_1 \wedge \neg Wt_2 \wedge \dots \wedge \neg Wt_{1T}) \quad (3)$$

But (3) is logically equivalent to the statement that there doesn't exist a winning ticket; i.e., (3) is logically equivalent to this result from Sequence 2:

$$\mathbf{B}_a \neg \exists t_i Wt_i \quad (4)$$

Sequence 2, “Rigorized”

4 As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$4 \quad \text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet.

Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:

$$\mathbf{B}_a^3 \neg Wt_1 \wedge \mathbf{B}_a^3 \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a^3 \neg Wt_{1T} \quad (2)$$

Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:

But (3) is logically equivalent to the statement that there doesn't exist a winning ticket; i.e., (3) is logically equivalent to this result from Sequence 2:

$$\mathbf{B}_a \neg \exists t_i Wt_i \quad (4)$$

Sequence 2, “Rigorized”

4 As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$4 \quad \text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet.

Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:

$$\mathbf{B}_a^3 \neg Wt_1 \wedge \mathbf{B}_a^3 \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a^3 \neg Wt_{1T} \quad (2)$$

Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:

$$\mathbf{B}_a^3 (\neg Wt_1 \wedge \neg Wt_2 \wedge \dots \wedge \neg Wt_{1T}) \quad (3)$$

But (3) is logically equivalent to the statement that there doesn't exist a winning ticket; i.e., (3) is logically equivalent to this result from Sequence 2:

$$\mathbf{B}_a \neg \exists t_i Wt_i \quad (4)$$

Sequence 2, “Rigorized”

4 As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$4 \quad \text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet.

Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:

$$\mathbf{B}_a^3 \neg Wt_1 \wedge \mathbf{B}_a^3 \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a^3 \neg Wt_{1T} \quad (2)$$

Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:

$$\mathbf{B}_a^3 (\neg Wt_1 \wedge \neg Wt_2 \wedge \dots \wedge \neg Wt_{1T}) \quad (3)$$

But (3) is logically equivalent to the statement that there doesn't exist a winning ticket; i.e., (3) is logically equivalent to this result from Sequence 2:

Sequence 2, “Rigorized”

4 As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$4 \quad \text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet.

Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:

$$\mathbf{B}_a^3 \neg Wt_1 \wedge \mathbf{B}_a^3 \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a^3 \neg Wt_{1T} \quad (2)$$

Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:

$$\mathbf{B}_a^3 (\neg Wt_1 \wedge \neg Wt_2 \wedge \dots \wedge \neg Wt_{1T}) \quad (3)$$

But (3) is logically equivalent to the statement that there doesn't exist a winning ticket; i.e., (3) is logically equivalent to this result from Sequence 2:

$$\mathbf{B}_a^3 \neg \exists t_i Wt_i \quad (4)$$

Paradox Solved!

Deduction preserves strength.

Clashes are resolved in favor of higher strength; clashes propagate backwards through inverse deduction; if no higher-strength factors, suspend belief.

Any proposition p such that $prob(p) < 1$ is at most evident.

Any rational belief that p , where the basis for p is at most evident, is at most an evident (= level 3) belief.

Paradox Solved!

Deduction preserves strength.

Clashes are resolved in favor of higher strength; clashes propagate backwards through inverse deduction; if no higher-strength factors, suspend belief.

$$B_a^4 \exists t_i W t_i \quad (3)$$

Any proposition p such that $\text{prob}(p) < 1$ is at most evident.

Any rational belief that p , where the basis for p is at most evident, is at most an evident (= level 3) belief.

Paradox Solved!

Deduction preserves strength.

Clashes are resolved in favor of higher strength; clashes propagate backwards through inverse deduction; if no higher-strength factors, suspend belief.

$$B_a^4 \exists t_i W t_i \quad (3)$$

$$B_a^3 \neg \exists t_i W t_i \quad (4)$$

Any proposition p such that $\text{prob}(p) < 1$ is at most evident.

Any rational belief that p , where the basis for p is at most evident, is at most an evident (= level 3) belief.

Paradox Solved!

Deduction preserves strength.

Clashes are resolved in favor of higher strength; clashes propagate backwards through inverse deduction; if no higher-strength factors, suspend belief.

$$B_a^4 \exists t_i W t_i \quad (3) \qquad B_a^3 \neg \exists t_i W t_i \quad (4)$$

Any proposition p such that $\text{prob}(p) < 1$ is at most evident.

Any rational belief that p , where the basis for p is at most evident, is at most an evident (= level 3) belief.

Paradox Solved!

Deduction preserves strength.

Clashes are resolved in favor of higher strength; clashes propagate backwards through inverse deduction; if no higher-strength factors, suspend belief.

$$\mathbb{B}_{\alpha}^{43} \ni t_i \ni W_i \ni t_i (3)(4)$$

Any proposition p such that $\text{prob}(p) < 1$ is at most evident.

Any rational belief that p , where the basis for p is at most evident, is at most an evident (= level 3) belief.

Paradox Solved!

Deduction preserves strength.

Clashes are resolved in favor of higher strength; clashes propagate backwards through inverse deduction; if no higher-strength factors, suspend belief.

$$B_a^4 \exists t_i W t_i \quad (3)$$

Any proposition p such that $\text{prob}(p) < 1$ is at most evident.

Any rational belief that p , where the basis for p is at most evident, is at most an evident (= level 3) belief.

Paradox Solved!

Deduction preserves strength.

Clashes are resolved in favor of higher strength; clashes propagate backwards through inverse deduction; if no higher-strength factors, suspend belief.

$$B_a^4 \exists t_i W t_i \quad (3)$$

Any proposition p such that $\text{prob}(p) < 1$ is at most evident.

Any rational belief that p , where the basis for p is at most evident, is at most an evident (= level 3) belief.

$$B_a^3 \neg W t_1 \wedge B_a^3 \neg W t_2 \wedge \dots \wedge B_a^3 \neg W t_{1T} \quad (2)$$

Paradox Solved!

Deduction preserves strength.

Clashes are resolved in favor of higher strength; clashes propagate backwards through inverse deduction; if no higher-strength factors, suspend belief.

$$B_a^4 \exists t_i W t_i \quad (3)$$

Any proposition p such that $\text{prob}(p) < 1$ is at most evident.

Any rational belief that p , where the basis for p is at most evident, is at most an evident (= level 3) belief.

$$B_a^3 \neg W t_1 \wedge B_a^3 \neg W t_2 \wedge \dots \wedge B_a^3 \neg W t_{1T} \quad (2)$$

Paradox Solved!

Deduction preserves strength.

Clashes are resolved in favor of higher strength; clashes propagate backwards through inverse deduction; if no higher-strength factors, suspend belief.

$$B_a^4 \exists t_i W t_i \quad (3)$$

Any proposition p such that $\text{prob}(p) < 1$ is at most evident.

Any rational belief that p , where the basis for p is at most evident, is at most an evident (= level 3) belief.

$$B_a^3 \neg W t_1 \wedge B_a^3 \neg W t_2 \wedge \dots \wedge B_a^3 \neg W t_{1T} \quad (2)$$

This is why, to Mega Millions ticket holder:
“Sorry. I’m rational, and I believe you won’t win.”

To be clear about the effects of the first principle:

$$\vdash \mathbf{B}_a^3 \neg \mathbf{X} x W x \wedge \mathbf{B}_a^3 \exists x W x!$$

$$\vdash \mathbf{B}_a^2 \neg \mathbf{X} x W x \wedge \mathbf{B}_a^2 \exists x W x!$$

$$\vdash \mathbf{B}_a^1 \neg \mathbf{X} x W x \wedge \mathbf{B}_a^1 \exists x W x!$$

Clashes are resolved in favor of higher strength; clashes propagate backwards through inverse deduction, preserving affirmation/belief of premises as far as is possible; if no higher-strength factors, suspend belief. (This means that in this case belief at level 4 also shoots down belief at level 2, and level 1. This is sort of bizarre, because to retain the belief (at levels 3, 2, 1) that every particular ticket won't win, the step that gets to believing the existential formula is blocked. Pollock doesn't have steps in his "arguments." Our agents thus ends up believing at all levels that some ticket will win, and believing at all levels 3 and down, of each particular ticket, that it won't win.)

slutten