

Gödel's Completeness Theorem

Selmer Bringsjord

Rensselaer AI & Reasoning (RAIR) Lab
Department of Cognitive Science
Department of Computer Science
Lally School of Management & Technology
Rensselaer Polytechnic Institute (RPI)
Troy, New York 12180 USA

Intro to Logic
4/22/2019



Gödel's Great Theorems (OUP)

by Selmer Bringsjord

- Introduction (“The Wager”)
- Brief Preliminaries (e.g. the propositional calculus)
- The Completeness Theorem
- The First Incompleteness Theorem
- The Second Incompleteness Theorem
- The Speedup Theorem
- The Continuum-Hypothesis Theorem
- The Time-Travel Theorem
- Gödel’s “God Theorem”
- Could a Machine Match Gödel’s Genius?



Gödel's Great Theorems (OUP)

by Selmer Bringsjord



- Introduction (“The Wager”)
- Brief Preliminaries (e.g. the propositional calculus)
- The Completeness Theorem
- The First Incompleteness Theorem
- The Second Incompleteness Theorem
- The Speedup Theorem
- The Continuum-Hypothesis Theorem
- The Time-Travel Theorem
- Gödel’s “God Theorem”
- Could a Machine Match Gödel’s Genius?



Gödel's Great Theorems (OUP)

by Selmer Bringsjord

- Introduction (“The Wager”)
- Brief Preliminaries (e.g. the propositional calculus)
- The Completeness Theorem
- The First Incompleteness Theorem
- The Second Incompleteness Theorem
- The Speedup Theorem
- The Continuum-Hypothesis Theorem
- The Time-Travel Theorem
- Gödel’s “God Theorem”
- Could a Machine Match Gödel’s Genius?



Gödel's Great Theorems (OUP)

by Selmer Bringsjord

- ➔ ● Introduction (“The Wager”)
- ➔ ● Brief Preliminaries (e.g. the propositional calculus)
- ➔ ● The Completeness Theorem
- The First Incompleteness Theorem
- The Second Incompleteness Theorem
- The Speedup Theorem
- The Continuum-Hypothesis Theorem
- The Time-Travel Theorem
- Gödel’s “God Theorem”
- Could a Machine Match Gödel’s Genius?



Some Timeline Points



Some Timeline Points

1906 Brünn, Austria-Hungary



Some Timeline Points

1923 Vienna

1906 Brünn, Austria-Hungary



Some Timeline Points

Undergrad in seminar by Schlick

1923 Vienna

1906 Brünn, Austria-Hungary



Some Timeline Points

1929 Doctoral Dissertation: Proof of Completeness Theorem
Undergrad in seminar by Schlick

1923 Vienna

1906 Brünn, Austria-Hungary



Some Timeline Points

1930 Announces (First) *Incompleteness* Theorem
1929 Doctoral Dissertation: Proof of *Completeness* Theorem
Undergrad in seminar by Schlick

1923 Vienna

1906 Brünn, Austria-Hungary



Some Timeline Points

1933 Hitler comes to power.

1930 Announces (First) *Incompleteness* Theorem

1929 Doctoral Dissertation: Proof of Completeness Theorem

Undergrad in seminar by Schlick

1923 Vienna

1906 Brünn, Austria-Hungary



Some Timeline Points

1940 Back to USA, for good.

1933 Hitler comes to power.

1930 Announces (First) *Incompleteness* Theorem

1929 Doctoral Dissertation: Proof of Completeness Theorem

Undergrad in seminar by Schlick

1923 Vienna

1906 Brünn, Austria-Hungary



Some Timeline Points



1940 Back to USA, for good.

1933 Hitler comes to power.

1930 Announces (First) *Incompleteness* Theorem

1929 Doctoral Dissertation: Proof of Completeness Theorem

Undergrad in seminar by Schlick

1923 Vienna

1906 Brünn, Austria-Hungary



Some Timeline Points

1978 Princeton NJ USA.



1940 Back to USA, for good.

1933 Hitler comes to power.

1930 Announces (First) *Incompleteness* Theorem

1929 Doctoral Dissertation: Proof of Completeness Theorem

Undergrad in seminar by Schlick

1923 Vienna

1906 Brünn, Austria-Hungary



Some Timeline Points

1978 Princeton NJ USA.



1940 Back to USA, for good.

1933 Hitler comes to power.

1930 Announces (First) *Incompleteness* Theorem

1929 Doctoral Dissertation: Proof of Completeness Theorem

Undergrad in seminar by Schlick

1923 Vienna

1906 Brünn, Austria-Hungary



Some Timeline Points

1978 Princeton NJ USA.



1940 Back to USA, for good.

1933 Hitler comes to power.

1930 Announces (First) *Incompleteness* Theorem

1929 Doctoral Dissertation: Proof of Completeness Theorem

Undergrad in seminar by Schlick

1923 Vienna

1906 Brünn, Austria-Hungary





R&W's Axiomatization of the Propositional Calculus

$$\text{A1} \quad (\phi \vee \phi) \rightarrow \phi$$

$$\text{A2} \quad \phi \rightarrow (\phi \vee \psi)$$

$$\text{A3} \quad (\phi \vee \psi) \rightarrow (\psi \vee \phi)$$

$$\text{A4} \quad (\psi \rightarrow \chi) \rightarrow ((\phi \vee \psi) \rightarrow (\phi \vee \chi))$$



R&W's Axiomatization of the Propositional Calculus

$$\text{A1} \quad (\phi \vee \phi) \rightarrow \phi$$

$$\text{A2} \quad \phi \rightarrow (\phi \vee \psi)$$

$$\text{A3} \quad (\phi \vee \psi) \rightarrow (\psi \vee \phi)$$

$$\text{A4} \quad (\psi \rightarrow \chi) \rightarrow ((\phi \vee \psi) \rightarrow (\phi \vee \chi))$$

All instances of these schemata are true no matter what the input (true or false). (Agreed?) And indeed every single formula in the propositional calculus that is true no matter what the permutation (as shown in a truth table), can be proved (somehow) from these four axioms (using the rules of inference given earlier in our semester). This, Gödel knew, and could use.

Completeness Theorem for The Propositional Calculus

Let Γ be a set $\{\phi_1, \phi_2, \dots\}$ of formulae in the the propositional calculus. Then either all of Γ are satisfiable, or the conjunction up to and including the point k (i.e. $\phi_1 \wedge \phi_2 \wedge \dots \wedge \phi_k$) of failure is refutable.

Completeness Theorem for The Propositional Calculus

Let Γ be a set $\{\phi_1, \phi_2, \dots\}$ of formulae in the the propositional calculus. Then either all of Γ are satisfiable, or the conjunction up to and including the point k (i.e. $\phi_1 \wedge \phi_2 \wedge \dots \wedge \phi_k$) of failure is refutable.

Completeness Theorem for The Propositional Calculus

Let Γ be a set $\{\phi_1, \phi_2, \dots\}$ of formulae in the the propositional calculus. Then either all of Γ are satisfiable, or the conjunction up to and including the point k (i.e. $\phi_1 \wedge \phi_2 \wedge \dots \wedge \phi_k$) of failure is refutable.

Let Γ be a set $\{\phi_1, \phi_2, \dots\}$ of formulae in the the propositional calculus. Then either all of Γ can be simultaneously true in some scenario, or the conjunction up to and including the point k (i.e. $\phi_1 \wedge \phi_2 \wedge \dots \wedge \phi_k$) of failure is **refutable** (i.e. $\vdash \neg(\phi_1 \wedge \phi_2 \wedge \dots \wedge \phi_k)$).

Completeness Theorem for The Propositional Calculus

Let Γ be a set $\{\phi_1, \phi_2, \dots\}$ of formulae in the the propositional calculus. Then either all of Γ are satisfiable, or the conjunction up to and including the point k (i.e. $\phi_1 \wedge \phi_2 \wedge \dots \wedge \phi_k$) of failure is refutable.

Let Γ be a set $\{\phi_1, \phi_2, \dots\}$ of formulae in the the propositional calculus. Then either all of Γ can be simultaneously true in some scenario, or the conjunction up to and including the point k (i.e. $\phi_1 \wedge \phi_2 \wedge \dots \wedge \phi_k$) of failure is **refutable** (i.e. $\vdash \neg(\phi_1 \wedge \phi_2 \wedge \dots \wedge \phi_k)$).

Show under doc cam that we can build the scenario if we don't get a closed branch (the scenario is an infinite branch), or we can use resolution/the Oracle_{PC} to obtain a contradiction once we have supposed for indirect proof that the conjunction holds.

But the assumption that
there *is* an infinite branch is
based on König's Lemma ...

Toward König's Lemma as Train Travel





“To infinity and beyond!”






König's Lemma (train-travel version)


In a one-way train-travel map with finitely many options leading from each station, if there are partial paths forward of every finite length, there is an *infinite* path (= a path “to infinity”).

























Exercise 2:

Is there an algorithm for traveling this way?

Exercise 2:

Is there an algorithm for traveling this way?

No. This strategy for travel is beyond the reach of standard computation.

Exercise 2:

Is there an algorithm for traveling this way?

No. This strategy for travel is beyond the reach of standard computation.

(Does it not then follow, assuming that humans can find and “use” a provably correct strategy for this travel, that humans can’t be fundamentally computing machines?)

Proving the Lemma

(that there is an infinite branch)

Proof: We are seeking to prove that there is an infinite path (= that you can keep going forward forever = that the number of your stops forward are the size of \mathbf{Z}^+).

To begin, assume the antecedent of the theorem (i.e. that, (1), there are finitely many options leading from each station, and that, (2), in the map there are partial paths forward of every finite size).

Now, you are standing at Penn Station (S_1), facing k options. At least one of these options must lead to partial paths of arbitrary size (the size of any m in \mathbf{Z}^+). (**Sub-Proof:** Suppose otherwise for indirect proof. Then there is some positive integer n that places a ceiling on the size of partial paths that can be reached. But this violates (2) — contradiction.) Proceed to choose one of these options that lead to partial paths of arbitrary size. You are now standing at a new station (S_2), one stop after Penn Station. At least one of these options must lead to partial parts of arbitrary size (the size of any m in \mathbf{Z}^+). (**Sub-Proof:** Suppose otherwise for indirect proof ...)

Since you can iterate this forever, you'll be on an infinite trip to infinity! Buzz will be happy.

Simple Buzz-Lightyear-Like Branch

<doc cam>

But how'd he do it for *FOL*??

But how'd he do it for *FOL*??

Arbitrary ϕ of \mathcal{L}_1 to $Q\psi$ to $\forall x\exists y\gamma(x, y)$ to $\gamma(a, b)$ to what we've seen!

For Further Reading

THE DISCOVERY OF MY COMPLETENESS PROOFS

LEON HENKIN

Dedicated to my teacher, Alonzo Church, in his 91st year.

§1. Introduction. This paper deals with aspects of my doctoral dissertation¹ which contributed to the early development of model theory. What was of use to later workers was less the results of my thesis, than the method by which I proved the completeness of first-order logic—a result established by Kurt Gödel in *his* doctoral thesis 18 years before.²

The ideas that fed my discovery of this proof were mostly those I found in the teachings and writings of Alonzo Church. This may seem curious, as his work in logic, and his teaching, gave great emphasis to the constructive character of mathematical logic, while the model theory to which I contributed is filled with theorems about very large classes of mathematical structures, whose proofs often by-pass constructive methods.

Another curious thing about my discovery of a new proof of Gödel's completeness theorem, is that it arrived in the midst of my efforts to prove an entirely different result. Such "accidental" discoveries arise in many parts of scientific work. Perhaps there are regularities in the conditions under which such "accidents" occur which would interest some historians, so I shall try to describe in some detail the accident which befell me.

Received November 17, 1995, and in revised form, January 4, 1996.

For Further Reading

THE DISCOVERY OF MY COMPLETENESS PROOFS

LEON HENKIN

Dedicated to my teacher, Alonzo Church, in his 91st year.

§1. Introduction. This paper deals with aspects of my doctoral dissertation¹ which contributed to the early development of model theory. What was of use to later workers was less the results of my thesis, than the method by which I proved the completeness of first-order logic—a result established by Kurt Gödel in *his* doctoral thesis 18 years before.²

The ideas that fed my discovery of this proof were mostly those I found in the teachings and writings of Alonzo Church. This may seem curious, as his work in logic, and his teaching, gave great emphasis to the constructive character of mathematical logic, while the model theory to which I contributed is filled with theorems about very large classes of mathematical structures, whose proofs often by-pass constructive methods.

Another curious thing about my discovery of a new proof of Gödel's completeness theorem, is that it arrived in the midst of my efforts to prove an entirely different result. Such "accidental" discoveries arise in many parts of scientific work. Perhaps there are regularities in the conditions under which such "accidents" occur which would interest some historians, so I shall try to describe in some detail the accident which befell me.

Received November 17, 1995, and in revised form, January 4, 1996.

For Further Reading

THE DISCOVERY OF MY COMPLETENESS PROOFS

LEON HENKIN

Dedicated to my teacher, Alonzo Church, in his 91st year.

§1. Introduction. This paper deals with aspects of my doctoral dissertation¹ which contributed to the early development of model theory. What was of use to later workers was less the results of my thesis, than the method by which I proved the completeness of first-order logic—a result established by Kurt Gödel in *his* doctoral thesis 18 years before.²

The ideas that fed my discovery of this proof were mostly those I found in the teachings and writings of Alonzo Church. This may seem curious, as his work in logic, and his teaching, gave great emphasis to the constructive character of mathematical logic, while the model theory to which I contributed is filled with theorems about very large classes of mathematical structures, whose proofs often by-pass constructive methods.

Another curious thing about my discovery of a new proof of Gödel's completeness theorem, is that it arrived in the midst of my efforts to prove an entirely different result. Such "accidental" discoveries arise in many parts of scientific work. Perhaps there are regularities in the conditions under which such "accidents" occur which would interest some historians, so I shall try to describe in some detail the accident which befell me.

Received November 17, 1995, and in revised form, January 4, 1996.

For Further Reading

THE DISCOVERY OF MY COMPLETENESS PROOFS

LEON HENKIN

Dedicated to my teacher, Alonzo Church, in his 91st year.

§1. Introduction. This paper deals with aspects of my doctoral dissertation¹ which contributed to the early development of model theory. What was of use to later workers was less the results of my thesis, than the method by which I proved the completeness of first-order logic—a result established by Kurt Gödel in *his* doctoral thesis 18 years before.²

The ideas that fed my discovery of this proof were mostly those I found in the teachings and writings of Alonzo Church. This may seem curious, as his work in logic, and his teaching, gave great emphasis to the constructive character of mathematical logic, while the model theory to which I contributed is filled with theorems about very large classes of mathematical structures, whose proofs often by-pass constructive methods.

Another curious thing about my discovery of a new proof of Gödel's completeness theorem, is that it arrived in the midst of my efforts to prove an entirely different result. Such "accidental" discoveries arise in many parts of scientific work. Perhaps there are regularities in the conditions under which such "accidents" occur which would interest some historians, so I shall try to describe in some detail the accident which befell me.

Received November 17, 1995, and in revised form, January 4, 1996.

For Further Reading

THE DISCOVERY OF MY COMPLETENESS PROOFS

LEON HENKIN

Dedicated to my teacher, Alonzo Church, in his 91st year.

§1. Introduction. This paper deals with aspects of my doctoral dissertation¹ which contributed to the early development of model theory. What was of use to later workers was less the results of my thesis, than the method by which I proved the completeness of first-order logic—a result established by Kurt Gödel in *his* doctoral thesis 18 years before.²

The ideas that fed my discovery of this proof were mostly those I found in the teachings and writings of Alonzo Church. This may seem curious, as his work in logic, and his teaching, gave great emphasis to the constructive character of mathematical logic, while the model theory to which I contributed is filled with theorems about very large classes of mathematical structures, whose proofs often by-pass constructive methods.

Another curious thing about my discovery of a new proof of Gödel's completeness theorem, is that it arrived in the midst of my efforts to prove an entirely different result. Such "accidental" discoveries arise in many parts of scientific work. Perhaps there are regularities in the conditions under which such "accidents" occur which would interest some historians, so I shall try to describe in some detail the accident which befell me.

Received November 17, 1995, and in revised form, January 4, 1996.

For Further Reading

THE DISCOVERY OF MY COMPLETENESS PROOFS

LEON HENKIN

Dedicated to my teacher, Alonzo Church, in his 91st year.

§1. Introduction. This paper deals with aspects of my doctoral dissertation¹ which contributed to the early development of model theory. What was of use to later workers was less the results of my thesis, than the method by which I proved the completeness of first-order logic—a result established by Kurt Gödel in *his* doctoral thesis 18 years before.²

The ideas that fed my discovery of this proof were mostly those I found in the teachings and writings of Alonzo Church. This may seem curious, as his work in logic, and his teaching, gave great emphasis to the constructive character of mathematical logic, while the model theory to which I contributed is filled with theorems about very large classes of mathematical structures, whose proofs often by-pass constructive methods.

Another curious thing about my discovery of a new proof of Gödel's completeness theorem, is that it arrived in the midst of my efforts to prove an entirely different result. Such "accidental" discoveries arise in many parts of scientific work. Perhaps there are regularities in the conditions under which such "accidents" occur which would interest some historians, so I shall try to describe in some detail the accident which befell me.

Received November 17, 1995, and in revised form, January 4, 1996.

slutten

Appendices ...

So what would be a *specific* g^* ? A truth of arithmetic that you can't move from the axioms of arithmetic?!?

So what would be a *specific* g^* ? A truth of arithmetic that you can't move from the axioms of arithmetic?!?

Here you go:

So what would be a *specific* g^* ? A truth of arithmetic that you can't move from the axioms of arithmetic?!?

Here you go:

That the Goodstein Sequence eventually reaches zero!

Goodstein Sequence; Goodstein's Theorem ...

Pure base n representation of a number r

- Represent r as only sum of powers of n in which the exponents are also powers of n etc

$$266 = 2^{2^{(2^{2^0} + 2^0)}} + 2^{(2^{2^0} + 2^0)} + 2^{2^0}$$

Grow Function

$Grow_k(n)$:

1. Take the pure base k representation of n
2. Replace all k by $k + 1$. Compute the number obtained.
3. Subtract one from the number

Example of **Grow**

$Grow_2(19)$

$$19 = 2^{2^{2^{2^0}}} + 2^{2^0} + 2^0$$

$$3^{3^{3^{3^0}}} + 3^{3^0} + 3^0$$

$$3^{3^{3^{3^0}}} + 3^{3^0} + 3^0 - 1$$

7625597484990

Goodstein Sequence

- For any natural number m

m

$Grow_2(m)$

$Grow_3(Grow_2(m))$

$Grow_4(Grow_3(Grow_2(m))),$

...

Sample Values

Sample Values

m										
2	2	2	1	0						

Sample Values

m										
2	2	2	1	0						
3	3	3	3	2	1	0				

Sample Values

m										
2	2	2	1	0						
3	3	3	3	2	1	0				
4	4	26	41	60	83	109	139	...	11327 (96th term)	...
5	15	$\sim 10^{13}$	$\sim 10^{155}$	$\sim 10^{2185}$	$\sim 10^{36306}$	10^{695975}	$10^{15151337}$...		

Yet, The Theorems!!

Yet, The Theorems!!

Theorem 1 (Goodstein's Theorem). *For all natural numbers, the Goodstein sequence reaches zero after a finite number of steps.*

Theorem 2 (Unprovability of Goodstein's Theorem). *Goodstein's theorem is not provable in Peano Arithmetic (PA) (or any equivalent theory of arithmetic).*

Yet, The Theorems!!

Theorem 1 (Goodstein's Theorem). *For all natural numbers, the Goodstein sequence reaches zero after a finite number of steps.*

Theorem 2 (Unprovability of Goodstein's Theorem). *Goodstein's theorem is not provable in Peano Arithmetic (PA) (or any equivalent theory of arithmetic).*

So, Gödel was right, empirically!

We have in GT a truth of elementary arithmetic that we can't prove from elementary arithmetic!

Could a computing machine get this?? ...

Small Steps Toward Hypercomputation via Infinitary Machine Proof Verification and Proof Generation

Naveen Sundar Govindarajulu, John Licato, and Selmer Bringsjord
Department of Computer Science
Department of Cognitive Science
Rensselaer AI & Reasoning Laboratory
govinn@rpi.edu • licatj@rpi.edu • selmer@rpi.edu

Rensselaer Polytechnic Institute
110 8th Street, Troy, NY 12180 USA

Abstract. After setting a context based on two general points (that humans appear to reason in infinitary fashion, and two, that actual hypercomputers aren't currently available to directly model and replicate such infinitary reasoning), we set a humble engineering goal of taking initial steps toward a computing machine that can reason in infinitary fashion. The initial steps consist in our outline of automated proof-verification and proof-discovery techniques for theorems independent of PA that seem to require an understanding and use of infinitary concepts. We specifically focus on proof-discovery techniques that make use of a marriage of analogical and deductive reasoning (which we call *analogico-deductive reasoning*).


A Context: Infinitary Reasoning, Hypercomputation, and Humble Engineering

Bringsjord has repeatedly pointed out the obvious fact that the behavior of formal scientists, taken at face value, involve various infinitary structures and reasoning. (We say "at face value" to simply indicate we don't presuppose some view that denies the reality of infinite entities routinely involved in the formal sciences.) For example, in (Bringsjord & van Heuveln 2003), Bringsjord himself operates as such a scientist in presenting an infinitary paradox which to his knowledge has yet to be solved. And he has argued that apparently infinitary behavior constitutes a grave challenge to AI and the Church-Turing Thesis (e.g., see Bringsjord & Arkoudas 2006, Bringsjord & Zenzen 2003). More generally, Bringsjord conjectures that every human-produced proof of a theorem independent of Peano Arithmetic (PA) will make use of infinitary structures and reasoning, when these structures are taken at face value¹. We have ourselves designed logico-computational logics for handling infinitary reasoning (e.g., see the treatment of the infinitized wise-man puzzle: Arkoudas & Bringsjord 2005), but this work simply falls back on the human ability to carry out induction on the natural numbers: it doesn't dissect and explain this ability. Finally, it must be admitted by all that there is simply no systematic, comprehensive model or framework anywhere in the formal/computational approach to understanding human knowledge and intelligence that provides a theory about how humans are able to engage with infinitary structures. This is revealed perhaps most clearly when one studies the fruit produced by the part of formal AI devoted to producing discovery systems: such fruit is embarrassingly finitary (e.g., see Shilliday 2009).


Given this context, we are interested in exploring how one might give a machine the ability to reason in infinitary fashion. We are not saying that we in fact have figured out how to give such ability to a computing machine. Our objective here is much more humble and limited: it is to push forward in the *attempt* to engineer a computing machine that has the ability to reason in infinitary fashion. Ultimately, if such an attempt is to succeed, the computing machine in question will presumably be capable of outright hypercomputation. But the fact is that from an engineering perspective, we don't know how to create and harness a hypercomputer. So what we must first try to do, as explained in (Bringsjord & Zenzen 2003), is pursue engineering that initiates the attempt to engineer a hypercomputer, and takes the first few steps. In the present paper, the engineering is aimed specifically at giving a computing machine the ability to, in a limited but well-defined sense, reason in infinitary fashion. Even more specifically, our engineering is aimed at building a machine capable of at least providing a strong case for a result which, in the human sphere, has hitherto required use of infinitary techniques.

¹ A weaker conjecture along the same line has been ventured by Isaacson, and is elegantly discussed by Smith (2007).

Needs Understanding of Ordinal Numbers ...




Needs Understanding of Ordinal Numbers ...



The image shows a window titled "Slate - Example.slt" with a white background. The window contains a mathematical inequality comparing two expressions. The left expression is $19_1 = 2^{2^{2^{2^0}}} + 2^{2^0} + 2^0$. The right expression is $\omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} + \omega^0$. The inequality is $19_1 = 2^{2^{2^{2^0}}} + 2^{2^0} + 2^0 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} + \omega^0$.

$$19_1 = 2^{2^{2^{2^0}}} + 2^{2^0} + 2^0 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} + \omega^0$$

Needs Understanding of Ordinal Numbers ...



Slate - Example.slt

$$19_1 = 2^{2^{2^{2^0}}} + 2^{2^0} + 2^0 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} + \omega^0$$
$$19_2 = 3^{3^{3^{3^0}}} + 3^{3^0} + 3^0 - 1 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} + \omega^0 - 1$$


Needs Understanding of Ordinal Numbers ...

$$19_1 = 2^{2^{2^{2^0}}} + 2^{2^0} + 2^0 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} + \omega^0$$

$$19_2 = 3^{3^{3^{3^0}}} + 3^{3^0} + 3^0 - 1 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} + \omega^0 - 1$$

$$19_3 = 4^{4^{4^{4^0}}} + 4^{4^0} - 1 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} - 1$$

Needs Understanding of Ordinal Numbers ...



Slate - Example.slt

$$19_1 = 2^{2^{2^{2^0}}} + 2^{2^0} + 2^0 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} + \omega^0$$
$$19_2 = 3^{3^{3^{3^0}}} + 3^{3^0} + 3^0 - 1 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} + \omega^0 - 1$$
$$19_3 = 4^{4^{4^{4^0}}} + 4^{4^0} - 1 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} - 1$$
$$19_4 = 5^{5^{5^{5^0}}} + 5^0 + 5^0 - 1 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^0 + \omega^0 - 1$$

Needs Understanding of Ordinal Numbers ...

Slate - Example.slt

$$19_1 = 2^{2^{2^{2^0}}} + 2^{2^0} + 2^0 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} + \omega^0$$
$$19_2 = 3^{3^{3^{3^0}}} + 3^{3^0} + 3^0 - 1 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} + \omega^0 - 1$$
$$19_3 = 4^{4^{4^{4^0}}} + 4^{4^0} - 1 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} - 1$$
$$19_4 = 5^{5^{5^{5^0}}} + 5^0 + 5^0 - 1 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^0 + \omega^0 - 1$$
$$19_5 = 6^{6^{6^{6^0}}} + 6^0 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^0$$

Needs Understanding of Ordinal Numbers ...

$$19_1 = 2^{2^{2^{2^0}}} + 2^{2^0} + 2^0 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} + \omega^0$$

$$19_2 = 3^{3^{3^{3^0}}} + 3^{3^0} + 3^0 - 1 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} + \omega^0 - 1$$

$$19_3 = 4^{4^{4^{4^0}}} + 4^{4^0} - 1 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} - 1$$

$$19_4 = 5^{5^{5^{5^0}}} + 5^0 + 5^0 - 1 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^0 + \omega^0 - 1$$

$$19_5 = 6^{6^{6^{6^0}}} + 6^0 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^0$$

⋮

Needs Understanding of Ordinal Numbers ...

Slate - Example.slt

$$19_1 = 2^{2^{2^{2^0}}} + 2^{2^0} + 2^0 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} + \omega^0$$
$$19_2 = 3^{3^{3^{3^0}}} + 3^{3^0} + 3^0 - 1 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} + \omega^0 - 1$$
$$19_3 = 4^{4^{4^{4^0}}} + 4^{4^0} - 1 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} - 1$$
$$19_4 = 5^{5^{5^{5^0}}} + 5^0 + 5^0 - 1 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^0 + \omega^0 - 1$$
$$19_5 = 6^{6^{6^{6^0}}} + 6^0 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^0$$

⋮

Needs Understanding of Ordinal Numbers ...

Slate - Example.slt

$$19_1 = 2^{2^{2^{2^0}}} + 2^{2^0} + 2^0 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} + \omega^0$$
$$19_2 = 3^{3^{3^{3^0}}} + 3^{3^0} + 3^0 - 1 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} + \omega^0 - 1$$
$$19_3 = 4^{4^{4^{4^0}}} + 4^{4^0} - 1 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} - 1$$
$$19_4 = 5^{5^{5^{5^0}}} + 5^0 + 5^0 - 1 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^0 + \omega^0 - 1$$
$$19_5 = 6^{6^{6^{6^0}}} + 6^0 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^0$$

⋮

strictly decreasing

$$19_1 = 2^{2^{2^{2^0}}} + 2^{2^0} + 2^0 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} + \omega^0$$

$$19_2 = 3^{3^{3^{3^0}}} + 3^{3^0} + 3^0 - 1 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} + \omega^0 - 1$$

$$19_3 = 4^{4^{4^{4^0}}} + 4^{4^0} - 1 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^{\omega^0} - 1$$


$$19_4 = 5^{5^{5^{5^0}}} + 5^0 + 5^0 - 1 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^0 + \omega^0 - 1$$

$$19_5 = 6^{6^{6^{6^0}}} + 6^0 < \omega^{\omega^{\omega^{\omega^0}}} + \omega^0$$

⋮

strictly decreasing

Ordinal Numbers ...



Yet, Conjecture (C)
(see “Isaacson’s Conjecture”)

Yet, Conjecture (**C**)

(see “Isaacson’s Conjecture”)

In order to produce a rationally compelling proof of any true sentence S formed from the symbol set of the language of arithmetic, but independent of **PA**, it’s necessary to deploy concepts and structures of an irreducibly infinitary nature.

Yet, Conjecture (**C**)

(see “Isaacson’s Conjecture”)

In order to produce a rationally compelling proof of any true sentence S formed from the symbol set of the language of arithmetic, but independent of **PA**, it’s necessary to deploy concepts and structures of an irreducibly infinitary nature.

If this is right, and computing machines can’t use irreducibly infinitary techniques, they’re in trouble — or: there won’t be a Singularity.