

The Lottery Paradox (and inductive logic)

Selmer Bringsjord

Rensselaer AI & Reasoning (RAIR) Lab
Department of Cognitive Science
Department of Computer Science
Lally School of Management & Technology
Rensselaer Polytechnic Institute (RPI)
Troy, New York 12180 USA

Intro to Logic
4/18/2019



R&D enabled by:



Paradoxes are engines of
progress in formal logic.

E.g., Russell's Paradox — as we've seen.

The paradox in this case?

The paradox in this case?

- A perfectly rational person can never believe P and believe $\neg P$ at the same time.

The paradox in this case?

- A perfectly rational person can never believe P and believe $\neg P$ at the same time.
- The Lottery Paradox (apparently) shows, courtesy of its two Sequences (of Reasoning), that a perfectly rational person can indeed have such a belief (upon considering a fair, large lottery).

The paradox in this case?

- A perfectly rational person can never believe P and believe $\neg P$ at the same time.
- The Lottery Paradox (apparently) shows, courtesy of its two Sequences (of Reasoning), that a perfectly rational person can indeed have such a belief (upon considering a fair, large lottery).
- Contradiction! — and hence a paradox!

Types of Paradoxes

- Deductive Paradoxes. The reasoning in question is exclusively deductive.
 - Russell's Paradox
 - The Liar Paradox
 - Richard's Paradox
- Inductive Paradoxes. Some of the reasoning in question uses non-deductive reasoning (e.g., probabilistic reasoning, abductive reasoning, analogical reasoning, etc.).

Types of Paradoxes

- Deductive Paradoxes. The reasoning in question is exclusively deductive.
 - Russell's Paradox
 - The Liar Paradox
 - Richard's Paradox
- Inductive Paradoxes Some of the reasoning in question uses non-deductive reasoning (e.g., probabilistic reasoning, abductive reasoning, analogical reasoning, etc.).

Types of Paradoxes

- Deductive Paradoxes. The reasoning in question is exclusively deductive.
 - Russell's Paradox
 - The Liar Paradox
 - Richard's Paradox
- Inductive Paradoxes. Some of the reasoning in question uses non-deductive reasoning (e.g., probabilistic reasoning, abductive reasoning, analogical reasoning, etc.).

Types of Paradoxes

- Deductive Paradoxes. The reasoning in question is exclusively deductive.
 - Russell's Paradox
 - The Liar Paradox
 - Richard's Paradox
- Inductive Paradoxes Some of the reasoning in question uses non-deductive reasoning (e.g., probabilistic reasoning, abductive reasoning, analogical reasoning, etc.).

Inductive Logic

the hallmark of deductive logic is *proof*, the hallmark of inductive logic is the concept of an *argument*. An exceptionally strong kind of argument is a proof, but plenty of arguments fall short of being proofs — and yet still have considerable force. For instance, consider the following argument α_1 :

- (1) Tweety is bird.
- (2) Most birds can fly.

- \therefore (3) Tweety can fly.

For stark contrast, consider as well this argument (α_2):

- (1') 3 is a positive integer.
- (2') All positive integers are greater than 0.

- \therefore (3') 3 is greater than 0.

The second of these arguments qualifies as an outright proof. That is, using the notation much employed before the present chapter:

$$\{(1'), (2')\} \vdash (3')$$

But in stark contrast, argument α_1 is not a proof that Tweety can fly. The reason is obvious: (3) isn't deduced from the combination of (1) and (2); that is,

$$\{(1), (2)\} \not\vdash (3)$$

Inductive Logic

the hallmark of deductive logic is *proof*, the hallmark of inductive logic is the concept of an *argument*. An exceptionally strong kind of argument is a proof, but plenty of arguments fall short of being proofs — and yet still have considerable force. For instance, consider the following argument α_1 :

$$\begin{array}{l} (1) \quad \text{Tweety is bird.} \\ (2) \quad \text{Most birds can fly.} \\ \hline \therefore (3) \quad \text{Tweety can fly.} \end{array}$$

For stark contrast, consider as well this argument (α_2):

$$\begin{array}{l} (1') \quad 3 \text{ is a positive integer.} \\ (2') \quad \text{All positive integers are greater than 0.} \\ \hline \therefore (3') \quad 3 \text{ is greater than 0.} \end{array}$$

The second of these arguments qualifies as an outright proof. That is, using the notation much employed before the present chapter:

$$\{(1'), (2')\} \vdash (3')$$

But in stark contrast, argument α_1 is not a proof that Tweety can fly. The reason is obvious: (3) isn't deduced from the combination of (1) and (2); that is,

$$\{(1), (2)\} \not\vdash (3)$$

Inductive Logic

the hallmark of deductive logic is *proof*, the hallmark of inductive logic is the concept of an *argument*. An exceptionally strong kind of argument is a proof, but plenty of arguments fall short of being proofs — and yet still have considerable force. For instance, consider the following argument α_1 :

- | | |
|------------------|---------------------|
| (1) | Tweety is bird. |
| (2) | Most birds can fly. |
| <hr/> | |
| \therefore (3) | Tweety can fly. |

For stark contrast, consider as well this argument (α_2):

- | | |
|-------------------|---|
| (1') | 3 is a positive integer. |
| (2') | All positive integers are greater than 0. |
| <hr/> | |
| \therefore (3') | 3 is greater than 0. |

Deductive (familiar)

The second of these arguments qualifies as an outright proof. That is, using the notation much employed before the present chapter:

$$\{(1'), (2')\} \vdash (3')$$

But in stark contrast, argument α_1 is not a proof that Tweety can fly. The reason is obvious: (3) isn't deduced from the combination of (1) and (2); that is,

$$\{(1), (2)\} \not\vdash (3)$$

Inductive Logic

the hallmark of deductive logic is *proof*, the hallmark of inductive logic is the concept of an *argument*. An exceptionally strong kind of argument is a proof, but plenty of arguments fall short of being proofs — and yet still have considerable force. For instance, consider the following argument α_1 :

(1)	Tweety is bird.
(2)	Most birds can fly.
<hr/>	
\therefore (3)	Tweety can fly.

For stark contrast, consider as well this argument (α_2):

(1')	3 is a positive integer.
(2')	All positive integers are greater than 0.
<hr/>	
\therefore (3')	3 is greater than 0.

Deductive (familiar)

The second of these arguments qualifies as an outright proof. That is, using the notation much employed before the present chapter:

$$\{(1'), (2')\} \vdash (3')$$

But in stark contrast, argument α_1 is not a proof that Tweety can fly. The reason is obvious: (3) isn't deduced from the combination of (1) and (2); that is,

$$\{(1), (2)\} \not\vdash (3)$$

Inductive Logic

the hallmark of deductive logic is *proof*, the hallmark of inductive logic is the concept of an *argument*. An exceptionally strong kind of argument is a proof, but plenty of arguments fall short of being proofs — and yet still have considerable force. For instance, consider the following argument α_1 :

(1)	Tweety is bird.
(2)	Most birds can fly.
<hr/>	
\therefore (3)	Tweety can fly.

Inductive (new)

For stark contrast, consider as well this argument (α_2):

(1')	3 is a positive integer.
(2')	All positive integers are greater than 0.
<hr/>	
\therefore (3')	3 is greater than 0.

Deductive (familiar)

The second of these arguments qualifies as an outright proof. That is, using the notation much employed before the present chapter:

$$\{(1'), (2')\} \vdash (3')$$

But in stark contrast, argument α_1 is not a proof that Tweety can fly. The reason is obvious: (3) isn't deduced from the combination of (1) and (2); that is,

$$\{(1), (2)\} \not\vdash (3)$$

Inductive-Reasoning Example from Pollock — for Peek Ahead

Imagine the following:

Keith tells you that the morning news predicts rain in Troy today. However, Alvin tells you that the same news report predicted sunshine.

Imagine the following: Keith tells you that the morning news predicts rain in Tucson today. However, Alvin tells you that the same news report predicted sunshine.

Without any other source of information, it would be irrational to place belief in either Keith's or Alvin's statements.

Imagine the following: Keith tells you that the morning news predicts rain in Tucson today. However, Alvin tells you that the same news report predicted sunshine.

Without any other source of information, it would be irrational to place belief in either Keith's or Alvin's statements.

Further, suppose you happened to watch the noon news report, and that report predicted rain. Then you should believe that it will rain despite your knowledge of Alvin's argument.

Defeasible Reasoning in OSCAR


K- Keith says that M

A- Alvin says that $\sim M$

M- The morning news said that R

R- It is going to rain this afternoon

N- The noon news says that R



All such can be absorbed into our inductive logics and our automated inductive reasoners (= our AI).

In Our Inductive Modal Logic

- | | | |
|-----|--|------|
| (1) | $\mathbf{K}(you, \mathbf{S}(keith, \mathbf{S}(m, rain)))$ | fact |
| (2) | $\mathbf{K}(you, \mathbf{S}(alvin, \mathbf{S}(m, \neg rain)))$ | fact |

In Our Inductive Modal Logic

	(1)	$\mathbf{K}(you, \mathbf{S}(keith, \mathbf{S}(m, rain)))$	fact
	(2)	$\mathbf{K}(you, \mathbf{S}(alvin, \mathbf{S}(m, \neg rain)))$	fact
\therefore	(3)	$\mathbf{S}(keith, \mathbf{S}(m, rain))$?
\therefore	(4)	$\mathbf{S}(alvin, \mathbf{S}(m, \neg rain))$?

In Our Inductive Modal Logic

	(1)	$\mathbf{K}(you, \mathbf{S}(keith, \mathbf{S}(m, rain)))$	fact
	(2)	$\mathbf{K}(you, \mathbf{S}(alvin, \mathbf{S}(m, \neg rain)))$	fact
\therefore	(3)	$\mathbf{S}(keith, \mathbf{S}(m, rain))$?
\therefore	(4)	$\mathbf{S}(alvin, \mathbf{S}(m, \neg rain))$?
	(5)	$\mathbf{S}(keith, \phi) \rightarrow \mathbf{B}^2(you, \phi)$	Testimonial P1

In Our Inductive Modal Logic

	(1)	$\mathbf{K}(you, \mathbf{S}(keith, \mathbf{S}(m, rain)))$	fact
	(2)	$\mathbf{K}(you, \mathbf{S}(alvin, \mathbf{S}(m, \neg rain)))$	fact
\therefore	(3)	$\mathbf{S}(keith, \mathbf{S}(m, rain))$?
\therefore	(4)	$\mathbf{S}(alvin, \mathbf{S}(m, \neg rain))$?
	(5)	$\mathbf{S}(keith, \phi) \rightarrow \mathbf{B}^2(you, \phi)$	Testimonial P1
\therefore	(6)	$\mathbf{B}^2(you, \mathbf{S}(m, rain)) \wedge \mathbf{B}^2(you, \mathbf{S}(m, \neg rain))$	

In Our Inductive Modal Logic

	(1)	$\mathbf{K}(you, \mathbf{S}(keith, \mathbf{S}(m, rain)))$	fact
	(2)	$\mathbf{K}(you, \mathbf{S}(alvin, \mathbf{S}(m, \neg rain)))$	fact
\therefore	(3)	$\mathbf{S}(keith, \mathbf{S}(m, rain))$?
\therefore	(4)	$\mathbf{S}(alvin, \mathbf{S}(m, \neg rain))$?
	(5)	$\mathbf{S}(keith, \phi) \rightarrow \mathbf{B}^2(you, \phi)$	Testimonial P1
\therefore	(6)	$\mathbf{B}^2(you, \mathbf{S}(m, rain)) \wedge \mathbf{B}^2(you, \mathbf{S}(m, \neg rain))$	
\therefore	(7)	$\neg \mathbf{B}^2(you, \mathbf{S}(m, rain)) \wedge \neg \mathbf{B}^2(you, \mathbf{S}(m, \neg rain))$	“Clash” Principle

In Our Inductive Modal Logic

(1)	$\mathbf{K}(you, \mathbf{S}(keith, \mathbf{S}(m, rain)))$	fact
(2)	$\mathbf{K}(you, \mathbf{S}(alvin, \mathbf{S}(m, \neg rain)))$	fact
\therefore	(3) $\mathbf{S}(keith, \mathbf{S}(m, rain))$?
\therefore	(4) $\mathbf{S}(alvin, \mathbf{S}(m, \neg rain))$?
(5)	$\mathbf{S}(keith, \phi) \rightarrow \mathbf{B}^2(you, \phi)$	Testimonial P1
\therefore	(6) $\mathbf{B}^2(you, \mathbf{S}(m, rain)) \wedge \mathbf{B}^2(you, \mathbf{S}(m, \neg rain))$	
\therefore	(7) $\neg \mathbf{B}^2(you, \mathbf{S}(m, rain)) \wedge \neg \mathbf{B}^2(you, \mathbf{S}(m, \neg rain))$	“Clash” Principle
(8)	$\mathbf{K}(you, \mathbf{S}(noonnews, rain))$	

In Our Inductive Modal Logic

(1)	$\mathbf{K}(you, \mathbf{S}(keith, \mathbf{S}(m, rain)))$	fact
(2)	$\mathbf{K}(you, \mathbf{S}(alvin, \mathbf{S}(m, \neg rain)))$	fact
\therefore	(3) $\mathbf{S}(keith, \mathbf{S}(m, rain))$?
\therefore	(4) $\mathbf{S}(alvin, \mathbf{S}(m, \neg rain))$?
(5)	$\mathbf{S}(keith, \phi) \rightarrow \mathbf{B}^2(you, \phi)$	Testimonial P1
\therefore	(6) $\mathbf{B}^2(you, \mathbf{S}(m, rain)) \wedge \mathbf{B}^2(you, \mathbf{S}(m, \neg rain))$	
\therefore	(7) $\neg \mathbf{B}^2(you, \mathbf{S}(m, rain)) \wedge \neg \mathbf{B}^2(you, \mathbf{S}(m, \neg rain))$	“Clash” Principle
(8)	$\mathbf{K}(you, \mathbf{S}(noonnews, rain))$	
\therefore	(9) $\mathbf{S}(noonnews, rain)$?

In Our Inductive Modal Logic

(1)	$\mathbf{K}(you, \mathbf{S}(keith, \mathbf{S}(m, rain)))$	fact
(2)	$\mathbf{K}(you, \mathbf{S}(alvin, \mathbf{S}(m, \neg rain)))$	fact
\therefore	(3) $\mathbf{S}(keith, \mathbf{S}(m, rain))$?
\therefore	(4) $\mathbf{S}(alvin, \mathbf{S}(m, \neg rain))$?
(5)	$\mathbf{S}(keith, \phi) \rightarrow \mathbf{B}^2(you, \phi)$	Testimonial P1
\therefore	(6) $\mathbf{B}^2(you, \mathbf{S}(m, rain)) \wedge \mathbf{B}^2(you, \mathbf{S}(m, \neg rain))$	
\therefore	(7) $\neg \mathbf{B}^2(you, \mathbf{S}(m, rain)) \wedge \neg \mathbf{B}^2(you, \mathbf{S}(m, \neg rain))$	“Clash” Principle
(8)	$\mathbf{K}(you, \mathbf{S}(noonnews, rain))$	
\therefore	(9) $\mathbf{S}(noonnews, rain)$?
(10)	$\mathbf{S}(noonnews, \phi) \rightarrow \mathbf{B}^3(you, \phi)$	Testimonial P2

In Our Inductive Modal Logic

(1)	$\mathbf{K}(you, \mathbf{S}(keith, \mathbf{S}(m, rain)))$	fact
(2)	$\mathbf{K}(you, \mathbf{S}(alvin, \mathbf{S}(m, \neg rain)))$	fact
\therefore	(3) $\mathbf{S}(keith, \mathbf{S}(m, rain))$?
\therefore	(4) $\mathbf{S}(alvin, \mathbf{S}(m, \neg rain))$?
(5)	$\mathbf{S}(keith, \phi) \rightarrow \mathbf{B}^2(you, \phi)$	Testimonial P1
\therefore	(6) $\mathbf{B}^2(you, \mathbf{S}(m, rain)) \wedge \mathbf{B}^2(you, \mathbf{S}(m, \neg rain))$	
\therefore	(7) $\neg \mathbf{B}^2(you, \mathbf{S}(m, rain)) \wedge \neg \mathbf{B}^2(you, \mathbf{S}(m, \neg rain))$	“Clash” Principle
(8)	$\mathbf{K}(you, \mathbf{S}(noonnews, rain))$	
\therefore	(9) $\mathbf{S}(noonnews, rain)$?
(10)	$\mathbf{S}(noonnews, \phi) \rightarrow \mathbf{B}^3(you, \phi)$	Testimonial P2
\therefore	(11) $\mathbf{B}^3(you, rain)$	

The Lottery Paradox ...





E: “Please go down to Stewart’s & get the T U.”



E: “Please go down to Stewart’s & get the T U.”








S: “I’m sorry, E, I’m afraid I can’t do that.
It would be irrational.”











E: “Please go down to Stewart’s & get the T U.”


S: “I’m sorry, E, I’m afraid I can’t do that.
It would be irrational.”

•••




Sequence I







Sequence I




Sequence I




Sequence I




Sequence I







Sequence 1




Sequence 2




Sequence 1




Sequence 2





Sequence 1




Sequence 2









Sequence 1

Sequence 2

Contradiction!

Sequence I

Sequence I

Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent a is fully apprised.

Sequence I

Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

Sequence I

Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

$$Wt_1 \oplus Wt_2 \oplus \dots \oplus Wt_{1T} \quad (1)$$

Sequence I

Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

$$Wt_1 \oplus Wt_2 \oplus \dots \oplus Wt_{1T} \quad (1)$$

We then deduce from this that there is at least one ticket that will win, a proposition represented — using standard notation — as:

Sequence I

Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

$$Wt_1 \oplus Wt_2 \oplus \dots \oplus Wt_{1T} \quad (1)$$

We then deduce from this that there is at least one ticket that will win, a proposition represented — using standard notation — as:

$$\exists t_i Wt_i \quad (2)$$

Sequence I

Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

$$Wt_1 \oplus Wt_2 \oplus \dots \oplus Wt_{1T} \quad (1)$$

We then deduce from this that there is at least one ticket that will win, a proposition represented — using standard notation — as:

$$\exists t_i Wt_i \quad (2)$$

Very well; perfectly clear so far. And now we can add another deductive step: Since our rational agent *a* can follow this deduction sequence to this point, and since **D** is assumed to be indubitable, it follows that our rational agent in turn believes (2); i.e., we conclude Sequence I by obtaining the following:

Sequence I

Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

$$Wt_1 \oplus Wt_2 \oplus \dots \oplus Wt_{1T} \quad (1)$$


We then deduce from this that there is at least one ticket that will win, a proposition represented — using standard notation — as:

$$\exists t_i Wt_i \quad (2)$$


Very well; perfectly clear so far. And now we can add another deductive step: Since our rational agent *a* can follow this deduction sequence to this point, and since **D** is assumed to be indubitable, it follows that our rational agent in turn believes (2); i.e., we conclude Sequence I by obtaining the following:

$$\mathbf{B}_a \exists t_i Wt_i \quad (3)$$

Sequence I



Sequence I



Sequence 2

Sequence 2

As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent a is fully apprised.

Sequence 2

As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using 'IT' to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

Sequence 2

As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using '1T' to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$\text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

Sequence 2

As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$\text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_i winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet. Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_i won't win sails through—and this of course works for each ticket. Hence we have:

Sequence 2

As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$\text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet. Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:

$$\mathbf{B}_a \neg Wt_1 \wedge \mathbf{B}_a \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a \neg Wt_{1T} \quad (2)$$

Sequence 2

As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$\text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet. Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:

$$\mathbf{B}_a \neg Wt_1 \wedge \mathbf{B}_a \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a \neg Wt_{1T} \quad (2)$$

Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:

Sequence 2

As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$\text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet. Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:

$$\mathbf{B}_a \neg Wt_1 \wedge \mathbf{B}_a \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a \neg Wt_{1T} \quad (2)$$

Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:

$$\mathbf{B}_a (\neg Wt_1 \wedge \neg Wt_2 \wedge \dots \wedge \neg Wt_{1T}) \quad (3)$$

Sequence 2

As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$\text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet. Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:

$$\mathbf{B}_a \neg Wt_1 \wedge \mathbf{B}_a \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a \neg Wt_{1T} \quad (2)$$

Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:

$$\mathbf{B}_a (\neg Wt_1 \wedge \neg Wt_2 \wedge \dots \wedge \neg Wt_{1T}) \quad (3)$$

But (3) is logically equivalent to the statement that there doesn't exist a winning ticket; i.e., (3) is logically equivalent to this result from Sequence 2:

Sequence 2

As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$\text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet. Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:


$$\mathbf{B}_a \neg Wt_1 \wedge \mathbf{B}_a \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a \neg Wt_{1T} \quad (2)$$

Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:


$$\mathbf{B}_a (\neg Wt_1 \wedge \neg Wt_2 \wedge \dots \wedge \neg Wt_{1T}) \quad (3)$$


But (3) is logically equivalent to the statement that there doesn't exist a winning ticket; i.e., (3) is logically equivalent to this result from Sequence 2:


$$\mathbf{B}_a \neg \exists t_i Wt_i \quad (4)$$




Sequence 2









$\checkmark \mathbf{B}_a \neg \exists t_i W t_i \quad \mathbf{B}_a \exists t_i W t_i \checkmark$
 (The contradiction we sketched earlier has arrived.)

A Solution to The Lottery Paradox ...

Strength-Factor Continuum

Certain

Improbable

Evidently False

Probable

Beyond Reasonable Belief

Certainly False

Counterbalanced

Evident

Beyond Reasonable Doubt

Strength-Factor Continuum

Certain

Evident

Beyond Reasonable Doubt

Probable

Counterbalanced

Improbable

Beyond Reasonable Belief

Evidently False

Certainly False

Actually, now ...

Actually, now ...

certain (6)
evident (5)
overwhelmingly likely (4)
beyond reasonable doubt (3)
likely (2)
more likely than not (1)
counterbalanced (0)

Actually, now ...

certain	(6)
evident	(5)
overwhelmingly likely	(4)
beyond reasonable doubt	(3)
likely	(2)
more likely than not	(1)
counterbalanced	(0)

... but let's use the simpler scheme.

Strength-Factor Continuum

Certain

Evident

Beyond Reasonable Doubt

Probable

Counterbalanced

Improbable

Beyond Reasonable Belief

Evidently False

Certainly False

Strength-Factor Continuum

Certain

Evident

Beyond Reasonable Doubt

Probable

..... Counterbalanced

Improbable

Beyond Reasonable Belief

Evidently False

Certainly False

Strength-Factor Continuum

Epistemically Positive

Certain

Evident

Beyond Reasonable Doubt

Probable

..... Counterbalanced

Improbable

Beyond Reasonable Belief

Evidently False

Certainly False

Strength-Factor Continuum

Epistemically Positive

Certain

Evident

Beyond Reasonable Doubt

Probable

..... Counterbalanced

Improbable

Beyond Reasonable Belief

Evidently False

Epistemically Negative

Certainly False

Strength-Factor Continuum

Epistemically Positive

Certain

Evident

Beyond Reasonable Doubt

Probable

.....

Counterbalanced

Improbable

Beyond Reasonable Belief

Evidently False

Epistemically Negative

Certainly False

Strength-Factor Continuum

Epistemically Positive

(4) Certain

(3) Evident

(2) Beyond Reasonable Doubt

(1) Probable

..... (0) Counterbalanced

(-1) Improbable

(-2) Beyond Reasonable Belief

(-3) Evidently False

Epistemically Negative

(-4) Certainly False

Strength-Factor Continuum

Epistemically Positive

(4) Certain

(3) Evident

(2) Beyond Reasonable Doubt

(1) Probable

..... (0) Counterbalanced

(-1) Improbable

(-2) Beyond Reasonable Belief

(-3) Evidently False

Epistemically Negative

(-4) Certainly False

Epistemically Positive

(4) Certain

(3) Evident

(2) Beyond Reasonable Doubt

(1) Probable

..... (0) Counterbalanced

(-1) Improbable

(-2) Beyond Reasonable Belief

(-3) Evidently False

Epistemically Negative

(-4) Certainly False

Key Principles

Epistemically Positive

(4) Certain

(3) Evident

(2) Beyond Reasonable Doubt

(1) Probable

..... (0) Counterbalanced

(-1) Improbable

(-2) Beyond Reasonable Belief

(-3) Evidently False

Epistemically Negative

(-4) Certainly False

Key Principles

Epistemically Positive

Deduction preserves strength.

(4) Certain

(3) Evident

(2) Beyond Reasonable Doubt

(1) Probable

..... (0) Counterbalanced

(-1) Improbable

(-2) Beyond Reasonable Belief

(-3) Evidently False

Epistemically Negative

(-4) Certainly False

Key Principles

Epistemically Positive

(4) Certain

Deduction preserves strength.

(3) Evident

Clashes are resolved in favor of higher strength.

(2) Beyond Reasonable Doubt

(1) Probable

..... (0) Counterbalanced

(-1) Improbable

(-2) Beyond Reasonable Belief

(-3) Evidently False

Epistemically Negative

(-4) Certainly False

Key Principles

Epistemically Positive

(4) Certain

Deduction preserves strength.

(3) Evident

Clashes are resolved in favor of higher strength.

(2) Beyond Reasonable Doubt

(1) Probable

..... **Any proposition p such that $\text{prob}(p) < 1$ is at most evident.**

(0) Counterbalanced

(-1) Improbable

(-2) Beyond Reasonable Belief

(-3) Evidently False

Epistemically Negative

(-4) Certainly False

Key Principles

Epistemically Positive

(4) Certain

Deduction preserves strength.

(3) Evident

Clashes are resolved in favor of higher strength.

(2) Beyond Reasonable Doubt

(1) Probable

..... **Any proposition p such that $\text{prob}(p) < 1$ is at most evident.**

(0) Counterbalanced

Any rational belief that p , where the basis for p is at most evident, is at most an evident (= level 3) belief.

(-1) Improbable

(-2) Beyond Reasonable Doubt Belief

(-3) Evidently False

Epistemically Negative

(-4) Certainly False

Sequence I, “Rigorized”

Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

$$Wt_1 \oplus Wt_2 \oplus \dots \oplus Wt_{1T} \quad (1)$$

We then deduce from this that there is at least one ticket that will win, a proposition represented — using standard notation — as:

$$\exists t_i Wt_i \quad (2)$$

Very well; perfectly clear so far. And now we can add another deductive step: Since our rational agent *a* can follow this deduction sequence to this point, and since **D** is assumed to be indubitable, it follows that our rational agent in turn believes (2); i.e., we conclude Sequence I by obtaining the following:

$$\mathbf{B}_a \exists t_i Wt_i \quad (3)$$

Sequence I, “Rigorized”

- 4 Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

$$Wt_1 \oplus Wt_2 \oplus \dots \oplus Wt_{1T} \quad (1)$$

We then deduce from this that there is at least one ticket that will win, a proposition represented — using standard notation — as:

$$\exists t_i Wt_i \quad (2)$$

Very well; perfectly clear so far. And now we can add another deductive step: Since our rational agent *a* can follow this deduction sequence to this point, and since **D** is assumed to be indubitable, it follows that our rational agent in turn believes (2); i.e., we conclude Sequence I by obtaining the following:

$$B_a \exists t_i Wt_i \quad (3)$$

Sequence I, “Rigorized”

- 4 Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

$$4 \quad Wt_1 \oplus Wt_2 \oplus \dots \oplus Wt_{1T} \quad (1)$$

We then deduce from this that there is at least one ticket that will win, a proposition represented — using standard notation — as:

$$\exists t_i Wt_i \quad (2)$$

Very well; perfectly clear so far. And now we can add another deductive step: Since our rational agent *a* can follow this deduction sequence to this point, and since **D** is assumed to be indubitable, it follows that our rational agent in turn believes (2); i.e., we conclude Sequence I by obtaining the following:

$$\mathbf{B}_a \exists t_i Wt_i \quad (3)$$

Sequence I, “Rigorized”

- 4 Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

$$4 \quad Wt_1 \oplus Wt_2 \oplus \dots \oplus Wt_{1T} \quad (1)$$

We then deduce from this that there is at least one ticket that will win, a proposition represented — using standard notation — as:

$$4 \quad \exists t_i Wt_i \quad (2)$$

Very well; perfectly clear so far. And now we can add another deductive step: Since our rational agent *a* can follow this deduction sequence to this point, and since **D** is assumed to be indubitable, it follows that our rational agent in turn believes (2); i.e., we conclude Sequence I by obtaining the following:

$$\mathbf{B}_a \exists t_i Wt_i \quad (3)$$

Sequence I, “Rigorized”

- 4 Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

$$4 \quad Wt_1 \oplus Wt_2 \oplus \dots \oplus Wt_{1T} \quad (1)$$

We then deduce from this that there is at least one ticket that will win, a proposition represented — using standard notation — as:

$$4 \quad \exists t_i Wt_i \quad (2)$$

Very well; perfectly clear so far. And now we can add another deductive step: Since our rational agent *a* can follow this deduction sequence to this point, and since **D** is assumed to be indubitable, it follows that our rational agent in turn believes (2); i.e., we conclude Sequence I by obtaining the following:

$$4 \quad B_a \exists t_i Wt_i \quad (3)$$

Sequence I, “Rigorized”

- 4 Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

$$4 \quad Wt_1 \oplus Wt_2 \oplus \dots \oplus Wt_{1T} \quad (1)$$

We then deduce from this that there is at least one ticket that will win, a proposition represented — using standard notation — as:

$$4 \quad \exists t_i Wt_i \quad (2)$$

Very well; perfectly clear so far. And now we can add another deductive step: Since our rational agent *a* can follow this deduction sequence to this point, and since **D** is assumed to be indubitable, it follows that our rational agent in turn believes (2); i.e., we conclude Sequence I by obtaining the following:

Sequence I, “Rigorized”

- 4 Let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that either ticket 1 will win or ticket 2 will win or ... or ticket 1,000,000,000,000 will win. Let's write this (exclusive) disjunction as follows:

$$4 \quad Wt_1 \oplus Wt_2 \oplus \dots \oplus Wt_{1T} \quad (1)$$

We then deduce from this that there is at least one ticket that will win, a proposition represented — using standard notation — as:

$$4 \quad \exists t_i Wt_i \quad (2)$$

Very well; perfectly clear so far. And now we can add another deductive step: Since our rational agent *a* can follow this deduction sequence to this point, and since **D** is assumed to be indubitable, it follows that our rational agent in turn believes (2); i.e., we conclude Sequence I by obtaining the following:

$$4 \quad B_a^4 \exists t_i Wt_i \quad (3)$$

Sequence 2, “Rigorized”

As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$\text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet.

Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:

$$\mathbf{B}_a \neg Wt_1 \wedge \mathbf{B}_a \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a \neg Wt_{1T} \quad (2)$$

Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:

$$\mathbf{B}_a (\neg Wt_1 \wedge \neg Wt_2 \wedge \dots \wedge \neg Wt_{1T}) \quad (3)$$

But (3) is logically equivalent to the statement that there doesn't exist a winning ticket; i.e., (3) is logically equivalent to this result from Sequence 2:

$$\mathbf{B}_a \neg \exists t_i Wt_i \quad (4)$$

Sequence 2, “Rigorized”

4

As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$\text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet. Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:

$$\mathbf{B}_a \neg Wt_1 \wedge \mathbf{B}_a \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a \neg Wt_{1T} \quad (2)$$

Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:

$$\mathbf{B}_a (\neg Wt_1 \wedge \neg Wt_2 \wedge \dots \wedge \neg Wt_{1T}) \quad (3)$$

But (3) is logically equivalent to the statement that there doesn't exist a winning ticket; i.e., (3) is logically equivalent to this result from Sequence 2:

$$\mathbf{B}_a \neg \exists t_i Wt_i \quad (4)$$

Sequence 2, “Rigorized”

4 As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$4 \quad \text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet. Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:

$$\mathbf{B}_a \neg Wt_1 \wedge \mathbf{B}_a \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a \neg Wt_{1T} \quad (2)$$

Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:

$$\mathbf{B}_a (\neg Wt_1 \wedge \neg Wt_2 \wedge \dots \wedge \neg Wt_{1T}) \quad (3)$$

But (3) is logically equivalent to the statement that there doesn't exist a winning ticket; i.e., (3) is logically equivalent to this result from Sequence 2:

$$\mathbf{B}_a \neg \exists t_i Wt_i \quad (4)$$

Sequence 2, “Rigorized”

4 As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$4 \quad \text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet.

Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:

Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:

$$\mathbf{B}_a(\neg Wt_1 \wedge \neg Wt_2 \wedge \dots \wedge \neg Wt_{1T}) \quad (3)$$

But (3) is logically equivalent to the statement that there doesn't exist a winning ticket; i.e., (3) is logically equivalent to this result from Sequence 2:

$$\mathbf{B}_a \neg \exists t_i Wt_i \quad (4)$$

Sequence 2, “Rigorized”

4 As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$4 \quad \text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet.

Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:

$$\mathbf{B}_a^3 \neg Wt_1 \wedge \mathbf{B}_a^3 \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a^3 \neg Wt_{1T} \quad (2)$$

Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:

$$\mathbf{B}_a(\neg Wt_1 \wedge \neg Wt_2 \wedge \dots \wedge \neg Wt_{1T}) \quad (3)$$

But (3) is logically equivalent to the statement that there doesn't exist a winning ticket; i.e., (3) is logically equivalent to this result from Sequence 2:

$$\mathbf{B}_a \neg \exists t_i Wt_i \quad (4)$$

Sequence 2, “Rigorized”

4 As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$4 \quad \text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet.

Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:

$$\mathbf{B}_a^3 \neg Wt_1 \wedge \mathbf{B}_a^3 \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a^3 \neg Wt_{1T} \quad (2)$$

Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:

But (3) is logically equivalent to the statement that there doesn't exist a winning ticket; i.e., (3) is logically equivalent to this result from Sequence 2:

$$\mathbf{B}_a \neg \exists t_i Wt_i \quad (4)$$

Sequence 2, “Rigorized”

4 As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$4 \quad \text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet.

Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:

$$\mathbf{B}_a^3 \neg Wt_1 \wedge \mathbf{B}_a^3 \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a^3 \neg Wt_{1T} \quad (2)$$

Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:

$$\mathbf{B}_a^3 (\neg Wt_1 \wedge \neg Wt_2 \wedge \dots \wedge \neg Wt_{1T}) \quad (3)$$

But (3) is logically equivalent to the statement that there doesn't exist a winning ticket; i.e., (3) is logically equivalent to this result from Sequence 2:

$$\mathbf{B}_a \neg \exists t_i Wt_i \quad (4)$$

Sequence 2, “Rigorized”

4 As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

4
$$\text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet.

Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:

$$\mathbf{B}_a^3 \neg Wt_1 \wedge \mathbf{B}_a^3 \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a^3 \neg Wt_{1T} \quad (2)$$

Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:

$$\mathbf{B}_a^3 (\neg Wt_1 \wedge \neg Wt_2 \wedge \dots \wedge \neg Wt_{1T}) \quad (3)$$

But (3) is logically equivalent to the statement that there doesn't exist a winning ticket; i.e., (3) is logically equivalent to this result from Sequence 2:

Sequence 2, “Rigorized”

4 As in Sequence 1, once again let **D** be a meticulous and perfectly accurate description of a 1,000,000,000,000-ticket lottery, of which rational agent *a* is fully apprised.

From **D** it obviously can be proved that the probability of a particular ticket t_i winning is 1 in 1,000,000,000,000. Using ‘1T’ to denote 1 trillion, we can write the probability for each ticket to win as a conjunction:

$$4 \quad \text{prob}(Wt_1) = \frac{1}{1,000,000,000,000} = \frac{1}{1T} \wedge \text{prob}(Wt_2) = \frac{1}{1T} \wedge \dots \wedge \text{prob}(Wt_{1T}) = \frac{1}{1T} \quad (1)$$

For the next step, note that the probability of ticket t_1 winning is lower than, say, the probability that if you walk outside a minute from now you will be flattened on the spot by a door from a 747 that falls from a jet of that type cruising at 35,000 feet. Since you have the rational belief that death won't ensue if you go outside (and have this belief precisely because you believe that the odds of your sudden demise in this manner are vanishingly small), the inference to the rational belief on the part of *a* that t_1 won't win sails through—and this of course works for each ticket. Hence we have:

$$\mathbf{B}_a^3 \neg Wt_1 \wedge \mathbf{B}_a^3 \neg Wt_2 \wedge \dots \wedge \mathbf{B}_a^3 \neg Wt_{1T} \quad (2)$$

Of course, if a rational agent believes *P*, and believes *Q* as well, it follows that that agent will believe the conjunction *P* & *Q*. Applying this principle to (2) yields:

$$\mathbf{B}_a^3 (\neg Wt_1 \wedge \neg Wt_2 \wedge \dots \wedge \neg Wt_{1T}) \quad (3)$$

But (3) is logically equivalent to the statement that there doesn't exist a winning ticket; i.e., (3) is logically equivalent to this result from Sequence 2:

$$\mathbf{B}_a^3 \neg \exists t_i Wt_i \quad (4)$$

Paradox Solved!

Deduction preserves strength.

Clashes are resolved in favor of higher strength; clashes propagate backwards through inverse deduction; if no higher-strength factors, suspend belief.

Any proposition p such that $prob(p) < 1$ is at most evident.

Any rational belief that p , where the basis for p is at most evident, is at most an evident (= level 3) belief.

Paradox Solved!

Deduction preserves strength.

Clashes are resolved in favor of higher strength; clashes propagate backwards through inverse deduction; if no higher-strength factors, suspend belief.

$$B_a^4 \exists t_i W t_i \quad (3)$$

Any proposition p such that $\text{prob}(p) < 1$ is at most evident.

Any rational belief that p , where the basis for p is at most evident, is at most an evident (= level 3) belief.

Paradox Solved!

Deduction preserves strength.

Clashes are resolved in favor of higher strength; clashes propagate backwards through inverse deduction; if no higher-strength factors, suspend belief.

$$B_a^4 \exists t_i W t_i \quad (3)$$

$$B_a^3 \neg \exists t_i W t_i \quad (4)$$

Any proposition p such that $\text{prob}(p) < 1$ is at most evident.

Any rational belief that p , where the basis for p is at most evident, is at most an evident (= level 3) belief.

Paradox Solved!

Deduction preserves strength.

Clashes are resolved in favor of higher strength; clashes propagate backwards through inverse deduction; if no higher-strength factors, suspend belief.

$$B_a^4 \exists t_i W t_i \quad (3) \qquad B_a^3 \neg \exists t_i W t_i \quad (4)$$

Any proposition p such that $\text{prob}(p) < 1$ is at most evident.

Any rational belief that p , where the basis for p is at most evident, is at most an evident (= level 3) belief.

Paradox Solved!

Deduction preserves strength.

Clashes are resolved in favor of higher strength; clashes propagate backwards through inverse deduction; if no higher-strength factors, suspend belief.

$$\mathbb{B}_{\alpha}^{43} \ni t_i \ni W_i \ni t_i (3)(4)$$

Any proposition p such that $\text{prob}(p) < 1$ is at most evident.

Any rational belief that p , where the basis for p is at most evident, is at most an evident (= level 3) belief.

Paradox Solved!

Deduction preserves strength.

Clashes are resolved in favor of higher strength; clashes propagate backwards through inverse deduction; if no higher-strength factors, suspend belief.

$$B_a^4 \exists t_i W t_i \quad (3)$$

Any proposition p such that $\text{prob}(p) < 1$ is at most evident.

Any rational belief that p , where the basis for p is at most evident, is at most an evident (= level 3) belief.

Paradox Solved!

Deduction preserves strength.

Clashes are resolved in favor of higher strength; clashes propagate backwards through inverse deduction; if no higher-strength factors, suspend belief.

$$B_a^4 \exists t_i W t_i \quad (3)$$

Any proposition p such that $\text{prob}(p) < 1$ is at most evident.

Any rational belief that p , where the basis for p is at most evident, is at most an evident (= level 3) belief.

$$B_a^3 \neg W t_1 \wedge B_a^3 \neg W t_2 \wedge \dots \wedge B_a^3 \neg W t_{1T} \quad (2)$$

Paradox Solved!

Deduction preserves strength.

Clashes are resolved in favor of higher strength; clashes propagate backwards through inverse deduction; if no higher-strength factors, suspend belief.

$$B_a^4 \exists t_i W t_i \quad (3)$$

Any proposition p such that $\text{prob}(p) < 1$ is at most evident.

Any rational belief that p , where the basis for p is at most evident, is at most an evident (= level 3) belief.

$$B_a^3 \neg W t_1 \wedge B_a^3 \neg W t_2 \wedge \dots \wedge B_a^3 \neg W t_{1T} \quad (2)$$

Paradox Solved!

Deduction preserves strength.

Clashes are resolved in favor of higher strength; clashes propagate backwards through inverse deduction; if no higher-strength factors, suspend belief.

$$B_a^4 \exists t_i W t_i \quad (3)$$

Any proposition p such that $\text{prob}(p) < 1$ is at most evident.

Any rational belief that p , where the basis for p is at most evident, is at most an evident (= level 3) belief.

$$B_a^3 \neg W t_1 \wedge B_a^3 \neg W t_2 \wedge \dots \wedge B_a^3 \neg W t_{1T} \quad (2)$$

This is why, to Mega Millions ticket holder:
“Sorry. I’m rational, and I believe you won’t win.”

slutten