#### **Selmer Bringsjord**

Rensselaer AI & Reasoning (RAIR) Lab
Department of Cognitive Science
Department of Computer Science
Lally School of Management & Technology
Rensselaer Polytechnic Institute (RPI)
Troy, New York 12180 USA



#### **Selmer Bringsjord**

Rensselaer AI & Reasoning (RAIR) Lab
Department of Cognitive Science
Department of Computer Science
Lally School of Management & Technology
Rensselaer Polytechnic Institute (RPI)
Troy, New York 12180 USA





#### **Selmer Bringsjord**

Rensselaer AI & Reasoning (RAIR) Lab
Department of Cognitive Science
Department of Computer Science
Lally School of Management & Technology
Rensselaer Polytechnic Institute (RPI)
Troy, New York 12180 USA







#### **Selmer Bringsjord**

Rensselaer AI & Reasoning (RAIR) Lab
Department of Cognitive Science
Department of Computer Science
Lally School of Management & Technology
Rensselaer Polytechnic Institute (RPI)
Troy, New York 12180 USA







#### **Selmer Bringsjord**

Rensselaer AI & Reasoning (RAIR) Lab
Department of Cognitive Science
Department of Computer Science
Lally School of Management & Technology
Rensselaer Polytechnic Institute (RPI)
Troy, New York 12180 USA

Note: This is a version
designed for those who have
had at least one universitylevel course in formal logic

with coverage through  $\mathcal{Z}_1$ .







# Background Context ...

- Introduction ("The Wager")
- Brief Preliminaries (e.g. the propositional calculus & FOL)
- The Completeness Theorem
- The First Incompleteness Theorem
- The Second Incompleteness Theorem
- The Speedup Theorem
- The Continuum-Hypothesis Theorem
- The Time-Travel Theorem
- Gödel's "God Theorem"
- Could a Finite Machine Match Gödel's Greatness?



- Introduction ("The Wager")
- Brief Preliminaries (e.g. the propositional calculus & FOL)
- The Completeness Theorem
- The First Incompleteness Theorem
- The Second Incompleteness Theorem
- The Speedup Theorem
- The Continuum-Hypothesis Theorem
- The Time-Travel Theorem
- Gödel's "God Theorem"
- Could a Finite Machine Match Gödel's Greatness?



- Introduction ("The Wager")
- Brief Preliminaries (e.g. the propositional calculus & FOL)
- The Completeness Theorem
- The First Incompleteness Theorem
- The Second Incompleteness
   Theorem
- The Speedup Theorem
- The Continuum-Hypothesis Theorem
- The Time-Travel Theorem
- Gödel's "God Theorem"
- Could a Finite Machine Match Gödel's Greatness?



- Introduction ("The Wager")
- Brief Preliminaries (e.g. the propositional calculus & FOL)
- The Completeness Theorem
- The First Incompleteness Theorem
- The Second Incompleteness Theorem
- The Speedup Theorem
- The Continuum-Hypothesis Theorem
- The Time-Travel Theorem
- Gödel's "God Theorem"
- Could a Machine Match Gödel's Genius?



by Selmer Bringsjord

- Introduction ("The Wager")
- Brief Preliminaries (e.g. the propositional calculus & FOL)
- The Completeness Theorem
- The First Incompleteness Theorem
- The Second Incompleteness Theorem
- The Speedup Theorem
- The Continuum-Hypothesis Theorem
- The Time-Travel Theorem
- Gödel's "God Theorem"
- Could a Machine Match Gödel's Genius?



By far the greatest of GGT; Selm's analysis based Sherlock Holmes' mystery "Silver Blaze."

 $\bar{P}$ : This sentence is unprovable.

 $\bar{P}$ : This sentence is unprovable.

Suppose that  $\bar{P}$  is true. Then we can immediately deduce that  $\bar{P}$  is provable, because here is a proof:  $\bar{P} \to \bar{P}$  is an easy theorem, and from it and our supposition we deduce  $\bar{P}$  by modus ponens. But since what  $\bar{P}$  says is that it's unprovable, we have deduced that  $\bar{P}$  is false under our initial supposition.

 $\bar{P}$ : This sentence is unprovable.

Suppose that  $\bar{P}$  is true. Then we can immediately deduce that  $\bar{P}$  is provable, because here is a proof:  $\bar{P} \to \bar{P}$  is an easy theorem, and from it and our supposition we deduce  $\bar{P}$  by modus ponens. But since what  $\bar{P}$  says is that it's unprovable, we have deduced that  $\bar{P}$  is false under our initial supposition.

Suppose on the other hand that  $\bar{P}$  is false. Then we can immediately deduce that  $\bar{P}$  is unprovable: Suppose for *reductio* that  $\bar{P}$  is provable; then  $\bar{P}$  holds as a result of some proof, but what  $\bar{P}$  says is that it's unprovable; and so we have contradiction. But since what  $\bar{P}$  says is that it's unprovable, and we have just proved that under our supposition, we arrive at the conclusion that  $\bar{P}$  is true.

 $\bar{P}$ : This sentence is unprovable.

Suppose that  $\bar{P}$  is true. Then we can immediately deduce that  $\bar{P}$  is provable, because here is a proof:  $\bar{P} \to \bar{P}$  is an easy theorem, and from it and our supposition we deduce  $\bar{P}$  by modus ponens. But since what  $\bar{P}$  says is that it's unprovable, we have deduced that  $\bar{P}$  is false under our initial supposition.

Suppose on the other hand that  $\bar{P}$  is false. Then we can immediately deduce that  $\bar{P}$  is unprovable: Suppose for *reductio* that  $\bar{P}$  is provable; then  $\bar{P}$  holds as a result of some proof, but what  $\bar{P}$  says is that it's unprovable; and so we have contradiction. But since what  $\bar{P}$  says is that it's unprovable, and we have just proved that under our supposition, we arrive at the conclusion that  $\bar{P}$  is true.

 $\mathsf{T}(\bar{P})$  iff (i.e., if & only if)  $\neg \mathsf{T}(\bar{P}) = \mathsf{F}(\bar{P})$ 

 $\bar{P}$ : This sentence is unprovable.

Suppose that  $\bar{P}$  is true. Then we can immediately deduce that  $\bar{P}$  is provable, because here is a proof:  $\bar{P} \to \bar{P}$  is an easy theorem, and from it and our supposition we deduce  $\bar{P}$  by modus ponens. But since what  $\bar{P}$  says is that it's unprovable, we have deduced that  $\bar{P}$  is false under our initial supposition.

Suppose on the other hand that  $\bar{P}$  is false. Then we can immediately deduce that  $\bar{P}$  is unprovable: Suppose for *reductio* that  $\bar{P}$  is provable; then  $\bar{P}$  holds as a result of some proof, but what  $\bar{P}$  says is that it's unprovable; and so we have contradiction. But since what  $\bar{P}$  says is that it's unprovable, and we have just proved that under our supposition, we arrive at the conclusion that  $\bar{P}$  is true.

 $\mathsf{T}(\bar{P})$  iff (i.e., if & only if)  $\neg \mathsf{T}(\bar{P}) = \mathsf{F}(\bar{P})$ Contradiction!

 $\dot{\bar{\pi}}=$  "I'm unprovable."

 $\bar{P}$ : This sentence is unprovable.

Suppose that  $\bar{P}$  is true. Then we can immediately deduce that  $\bar{P}$  is provable, because here is a proof:  $\bar{P} \to \bar{P}$  is an easy theorem, and from it and our supposition we deduce  $\bar{P}$  by modus ponens. But since what  $\bar{P}$  says is that it's unprovable, we have deduced that  $\bar{P}$  is false under our initial supposition.

Suppose on the other hand that  $\bar{P}$  is false. Then we can immediately deduce that  $\bar{P}$  is unprovable: Suppose for *reductio* that  $\bar{P}$  is provable; then  $\bar{P}$  holds as a result of some proof, but what  $\bar{P}$  says is that it's unprovable; and so we have contradiction. But since what  $\bar{P}$  says is that it's unprovable, and we have just proved that under our supposition, we arrive at the conclusion that  $\bar{P}$  is true.

 $\mathsf{T}(\bar{P})$  iff (i.e., if & only if)  $\neg \mathsf{T}(\bar{P}) = \mathsf{F}(\bar{P})$ Contradiction!

All of this is fishy; but Gödel, as we've seen, transformed it (by e.g. use of his encryption scheme) into utterly precise, impactful, indisputable reasoning ...


#### PA (Peano Arithmetic):


A1 
$$\forall x(0 \neq s(x))$$
  
A2  $\forall x \forall y(s(x) = s(y) \rightarrow x = y)$   
A3  $\forall x(x \neq 0 \rightarrow \exists y(x = s(y)))$   
A4  $\forall x(x + 0 = x)$   
A5  $\forall x \forall y(x + s(y) = s(x + y))$   
A6  $\forall x(x \times 0 = 0)$   
A7  $\forall x \forall y(x \times s(y) = (x \times y) + x)$ 


And, every sentence that is the universal closure of an instance of


$$([\phi(0) \land \forall x(\phi(x) \to \phi(s(x)))] \to \forall x\phi(x))$$

where  $\phi(x)$  is open wff with variable x, and perhaps others, free.









"We can't use math to ascertain whether mathematics is consistent."

"If we are restricted to certain kinds of formal reasoning, and feel we must have all of **PA** (math, engineering, etc.), we can't ascertain whether mathematics is consistent."

Suppose  $\Phi \supset \mathbf{PA}$  that is

- (i) Con  $\Phi$ ;
- (ii) Turing-decidable (i.e. membership in  $\Phi$  is Turing-decidable); and
- (iii) sufficiently expressive to capture all of the operations of a Turing machine (i.e. Repr  $\Phi$ ).

Then  $\Phi \not\vdash \text{consis}_{\Phi}$ .

Suppose  $\Phi \supset \mathbf{PA}$  that is

Remember Church's Theorem!

- (i) Con  $\Phi$ ;
- (ii) Turing-decidable (i.e. membership in  $\Phi$  is Turing-decidable); and
- (iii) sufficiently expressive to capture all of the operations of a Turing machine (i.e. Repr  $\Phi$ ).

Then  $\Phi \not\vdash \text{consis}_{\Phi}$ .

Suppose  $\Phi \supset \mathbf{PA}$  that is

Remember Church's Theorem!

- (i) Con  $\Phi$ ;
- (ii) Turing-decidable (i.e. membership in  $\Phi$  is Turing-decidable); and
- (iii) sufficiently expressive to capture all of the operations of a Turing machine (i.e. Repr  $\Phi$ ).

Then  $\Phi \not\vdash \text{consis}_{\Phi}$ .

# To prove G2, we shall once again allow ourselves ...


#### The Fixed Point Theorem (FPT)

Assume that  $\Phi$  is a set of arithmetic sentences such that Repr  $\Phi$ . There for every arithmetic formula  $\psi(x)$  with one free variable x, there is an arithmetic sentence  $\phi$  s.t.

$$\Phi \vdash \phi \leftrightarrow \psi(\hat{n}^{\phi}).$$

We can intuitively understand  $\phi$  to be saying: "I have the property ascribed to me by the formula  $\psi$ ."

# FPT in HyperSlate®!



Ok; so let's do it ... and let's see if you can see why Gödel declared G2 to be a direct "corollary" of GI, and didn't bother to prove it in his original paper ...

**Proof**: Suppose that the antecedent (i)—(iii) of **G2** holds. Suppose for reductio that

$$\Phi \vdash \mathsf{consis}_{\Phi}$$
.

We need three ingredients, and we shall be done. First, from FPT we can again directly obtain:

(\*) 
$$\Phi \vdash \mathcal{G} \leftrightarrow \neg \mathcal{P}_{\Phi}(\hat{n}^{\mathcal{G}'})$$
.

Next, we can prove (how? ... from one half of GI!) that:

(7.9) If Con 
$$\Phi$$
, then  $\Phi \not\vdash \mathcal{G}$ .

Thirdly, we can logicize the meta-logical proposition that  $\Phi$  is consistent as an object-level conditional which can itself be proved formally from  $\Phi$ :

(\*\*) 
$$\Phi \vdash \operatorname{consis}_{\Phi} \to \neg \mathscr{P}_{\Phi}(\hat{n}^{\mathscr{G}}).$$

Contradiction! (Can you find it?) **QED** 

# Med nok penger, kan logikk løse alle problemer.